



GLYCEMIC CONTROL IS ONE OF THE PREVENTIVE MEASUREMENT FROM RETINOPATHY IN THE PATIENTS OF TYPE –I DIABETES MELLITUS

Rafiq Ahmed Jalbani¹, RoohiNaz¹, Ali Raza Memon^{1*}, Zainab Memon² and Ashok Narsani³

¹Department of Biochemistry, Liaquat University of Medical & Health Sciences (LUMHS) Jamshoro, Sindh, Pakistan.

²Multidisciplinary Research Laboratory at Medical Research Centre (MRC), Liaquat University of Medical & Health Sciences (LUMHS) Jamshoro, Sindh, Pakistan.

³Department of Ophthalmology, Eye Hospital Liaquat University of Medical & Health Sciences (LUMHS) Jamshoro, Sindh, Pakistan.

Corresponding Author:- **Ali Raza Memon**

E-mail: aliraza.biolumhs@yahoo.com

Article Info	ABSTRACT
<p>Received 25/04/2014 Revised 18/05/2014 Accepted 20/05/2014</p> <p>Key words: Type - I Diabetes mellitus, Diabetic Retinopathy, HbA1c.</p>	<p>Diabetic Retinopathy is one of the main complication of both types; type I & II Diabetes mellitus. HbA1c is the gold standard parameter to estimate the glycemic index nowadays. Poor glycemic index means high blood sugar level which is the proceeding factor for development of retinopathy. Total 80 subjects were include in the study, from which 40 were type - I diabetes without retinopathy and 40 were diabetes type - I with retinopathy. HbA1c was estimated by Bio Red Variant. Retinopathy was assessed by indirect ophthalmoscope. The mean of HbA1c Type - I diabetic patients without retinopathy was $9.87 \pm 0.85\%$ and in Type - I diabetes mellitus with retinopathy as, $11.27 \pm 0.77\%$ respectively. These results showed that glycemic control in type - I diabetes with retinopathy highly poor than type - I diabetic patients without retinopathy. Poor glycemic index take a part in proceeding diabetic complications like retinopathy.</p>

INTRODUCTION

According to the World Health Organization, in 2000 at least 171 million people worldwide suffered from diabetes mellitus, or 2.8% of the population of all over the world [1]. In 2007, 246 million people world-wide suffered from diabetes making the disease one of the most common non-communicable global diseases and the fourth leading cause of death in the world according to IDF estimates [2]. Diabetes mellitus occurs throughout the world, but is more common (especially type II) in the more developed countries. The greatest increase in prevalence is, however, expected to occur in Asia and Africa, where most patients will probably be found up to 2030 [1]. Pakistan is one of the South Asian countries with population of 150 millions

❖ Enhanced glucose flux through the polyol pathway

and prevalence of Diabetes in Pakistan is high ranging from 7.6% to 11% of adult population[3, 4].

Diabetic Retinopathy is the damage to the retina caused by complications of diabetes mellitus, which can eventually lead to blindness. It is an ocular manifestation of systemic disease which affects up to 80% of all patients who have had diabetes for 10 years or more[5].

The incidence of diabetic retinopathy in Pakistan is 9.1% to 13% of diabetic population[6]. Currently 4 major biochemical pathways have been hypothesized to explain the mechanism of diabetic eye diseases; all starting initially from hyperglycemia induced vascular injury[7].

These mainly include:



- ❖ Increased intracellular formation of advanced glycation end products
- ❖ Activation of protein kinase C isoforms
- ❖ Stimulation of hexosamine pathway

HbA_{1c} is the major fraction, constituting approximately 80% of Hb_{A1}. It is formed by the condensation of glucose with the N-terminal valine residue of each beta chain of HbA to form an unstable Schiff base (aldimine). The Schiff base may either dissociate or undergo an amadori rearrangement to form a stable ketoamine[8].

Nowadays normally cut point of HbA_{1c} is 6.5% of total Hb in non-diabetic individuals [9].

The risk of diabetic retinopathy increases stepwise with increasing degrees of hyperglycemia [10].

Many epidemiological studies have demonstrated a strong relation between hyperglycemia and the development and progression of diabetic retinopathy. Elevated HbA_{1c} levels have been found to be a potent predictor of progression of proliferative diabetic retinopathy. The interaction of advanced glycation end products and their receptors and increased activity of the polyol pathway have been implicated as mediators of increased micro vascular permeability, ischemia and angiogenesis [11].

MATERIAL & METHODS

The current study was carried out at the Department of Biochemistry with collaboration of Department of Allied Medicine & Department of Ophthalmology of Liaquat University of Medical & Health Sciences Jamshoro Sindh Pakistan. Total 80 patients were enrolled from them 40 were diagnosed cases of Type - I

diabetes mellitus without retinopathy & 40 were diagnosed cases of Type - I diabetes mellitus with retinopathy.

Sample Preparation

2ml of blood sample drawn by venipuncture under aseptic measure from each subject under study then transferred the blood into anticoagulant bottle containing EDTA. Samples were first diluted with hemolysis reagent then incubated at 18 -28 °C for minimum of 30 minutes.

Analytic Method

HbA_{1c} level will be measured by BIO-RED VARIANT™ HemoglobinA_{1c} program, utilized the principle of ion – exchange high performance liquid chromatography (HPLC) for automatic and accurate separation of HbA_{1c}. Retinopathy was assist by indirect ophthalmoscope.

Statistical Analysis

SPSS version 16 was applied to show the significance of the above study. Independent student t- test was applied to observe the significance value (p.value) between these two groups. MS word & Excel 2007 used or generate the tables & graphs.

RESULTS


The mean fasting blood glucose levels in type - I diabetes mellitus without retinopathy was 127.46 ± 11.45 mg/dl and in retinopathy it was 201.90 ± 10.65 mg/dl (p.value<0.01) where the mean HbA_{1c} level in diabetic patients without retinopathy was 9.87 ± 0.85 % while in diabetic patients with retinopathy it was 11.27 ± 0.77 % (p.value< 0.05) these results were shown in following table and graphs.

Table 1. Significance relation of poor glycemic index in development retinopathy


Variable	Type - I Diabetes Mellitus without Retinopathy	Type - I Diabetes mellitus with Retinopathy.
FBS (mg/dl)	127.46 ± 11.45	201.90 ± 10.65
HbA _{1c} %	9.87 ± 0.85	11.27 ± 0.77

The above results show that there is significance relation of poor glycemic index in development retinopathy.

Graph 1. Fasting Blood Sugar Levels



Graph 2. HbA_{1c} Levels



DISCUSSION

HbA_{1c} is the gold standard parameter to rule out the blood glucose level status in diabetic patients. Poor glycemic control significantly observed in different in different complications of diabetes mellitus as in diabetic retinopathy. Different studies regarding this problem also has been done like: Santos et al. (2005) [10] have done cross sectional study on 210 patients from them 99 patients of diabetes mellitus without retinopathy and 111 patients of diabetes mellitus with retinopathy. They demonstrated that glycemic control (HbA_{1c}) were more poor in diabetic retinopathy (p<0.001).

Farhan et al. (2005) [13] have reported in their study that poor glycemic control (p<0.0001) and longer duration of diabetes mellitus (p<0.0001) were important predicting factors of diabetic retinopathy.

Rama and Pradeepa 2007, [14] Ishrat Kareem et al. 2004 [15], Zélia Maria da Silva Corrêa1 et al [16] also reported in their studies that poor glycemic control and

longer duration of diabetes mellitus were more prone to development of diabetic complications like retinopathy.

The present study shows that the levels of HbA_{1c} % are significantly raised in type –II diabetic patients with retinopathy as compare with type- II diabetic patients without retinopathy (p<0.001). This concluded that good glycemic control can prevent the diabetic patients from retinopathy, so regular monitoring of fasting blood sugar level (FBS) and HbA_{1c} % must necessary for diabetic patients from prevention of diabetic complication like retinopathy.

CONCLUSION

From the current study it has been concluded that good glycemic control can prevent the diabetic patients from retinopathy, so regular monitoring of fasting blood sugar level (FBS) and HbA_{1c} % must necessary for diabetic patients from prevention of diabetic complication like retinopathy.

REFERENCES

1. Wild S, Roglic G, Green A, Sicree R, King H. (2004). Global prevalence of diabetes, estimates for 2000 and projections for 2030. *Diabetes Care*, 27(5), 1047–53.
2. Samreen Riaz. (2009). Diabetes Mellitus. *Scientific Research and Essay*, 4(5), 367-373.
3. International diabetes federation. (2010). Prevalence estimation D.M 2010 MENA IDF Diabetes Atlas.
4. Jaffar TH, Levcy AS, White FM et al. (2004). Ethnic differ & determinants of diabetes & central obesity among south Asians of Pakistan. *Diabetic Medicine*, 21(7), 716–723.
5. Kertes PJ, Johnson TM. (2007). Evidence Based Eye Care. In *Evidence Base for Diabetic Care*, by Jhon Willey & sons. 30th edition, 501 -511.
6. Kazi AA, Laghari NA, Jatoi SM. (2005). Current Trends in Diabetic Retinopathy. *JLUMHS*, 1-2.
7. Balasuramanyam M, Rema M, Premanand C. (2002). Biochemical and molecular mechanisms of diabetic retinopathy. *Curr Sci*, 83, 1506.
8. Sacks DB. (2002). Carbohydrates. *Text book of Clinical Chemistry*. 4th edition, Philadelphia, W.B Saunders, 878 - 884.
9. World Health Organization Report. (2011). Use of HbA_{1c} in the diagnosis of Diabetes Mellitus, 6-7.
10. Santos KG, Tschiede B, Schneider JR, Souto KEP, Roisenberg I. (2005). Prevalence of retinopathy in Caucasian type 2 diabetic patients from the south of Brazil and relationship with clinical and metabolic factors. *Braz J Med Res*, 38 (2), 221-225.
11. Kotoula. Maria G, Koukoulis. George, Zintzaras, Karabatsas, Constantions H, Chatzoulis, Dimitrios Z. (2005). Metabolic control of diabetes is associated with an improved response of diabetic retinopathy to panretinal photocoagulation. *Diabetes Care*, 28, 2454-2457.
12. Nathan DM, Kuenen J, Borg R, Zheug H, Schoenfeld, Heene RJ. (2008). Translating the A1C assays into estimate average glucose values *Diabetic care*, 31(8), 1473-1478.
13. Al-Shammari FKH, A1-Meraghi O, Nasif A, Al-Otaibi S. (2005). The prevalence of diabetic retinopathy and associated risk factors in type II diabetes mellitus in al-Naeem area (Kuwait). *Middle East Journal of Family Medicine*, 3(2), 1–8.
14. Rama M, Pradeepa R. (2007). Diabetic Retinopathy, an Indian perspective. *Indian J Med Res*, 125, 297-310.
15. Kareem I, Jaweed SA, Bardapurka JS, Patil VP. (2004). Study of magnesium, glycosylated hemoglobin, lipid profile in diabetic retinopathy. *Indian journal of Clinical Biochemistry*, 19 (2), 124-127.
16. Zélia Maria da Silva Corrêa1, André Moraes Freitas, Italo Mundialino Marcon. (2003). Risk factors related to the severity of diabetic retinopathy. *Arq Bras Oftimol*, 66, 739-743.

