



## EFFECT OF CAFFEINE ON PHYSICAL PERFORMANCE IN YOUNG ADULTS

Gaikwad MA<sup>1</sup>, Rawekar AT<sup>2</sup>, Waghmare L<sup>2</sup> and Biswas DA<sup>3</sup>

<sup>1</sup>Jr-3 Dept. of Physiology, JNMC, Sawangi (M), Wardha, Maharashtra, India.

<sup>2</sup>Asso.Prof. Dept. of Physiology, JNMC, Sawangi (M), Wardha, Maharashtra, India.

<sup>3</sup>Prof & HOD Dept. of Physiology, JNMC, Sawangi (M), Wardha, Maharashtra, India.

Corresponding Author:- Gaikwad MA

E-mail: [megha.gaikwad29@gmail.com](mailto:megha.gaikwad29@gmail.com)

### Article Info

Received 25/05/2014

Revised 19/06/2014

Accepted 22/06/2014

**Key words:** Caffeine,  
Physical  
Performance,  
Exercise Intensity,  
Muscle Strength,  
Adults.

### ABSTRACT

Caffeine (1,3,7-trimethylxanthine) is a habitual substance present in a wide variety of beverages and in chocolate-based foods and it is also used as adjuvant in some drugs. There have been numerous reports that caffeine is an ergogenic aid; ingestion of the drug has been shown to increase endurance, particularly in prolonged exercise lasting 30–120 min. The present study was undertaken to study the effects of caffeine in coffee on endurance exercise in young healthy adult males. Twenty young males participated in this single blind (caffeine 5mg·kg<sup>-1</sup> body weight) study. Participants underwent two testing sessions separated by 7 days, consisting of handgrip strength test, Rate of Perceived Exertion scale (RPE) test and an incremental test to exhaustion on treadmill. Paired 't' test revealed significant difference in handgrip strength test, RPE scale, exhaustion time, maximal oxygen uptake (VO<sub>2</sub>max) and heart rate ( $p > 0.05$ ) before and after caffeine ingestion. The present study indicates that caffeine consumed 60 minutes before exercise can enhance exercise performance by increasing the total time to fatigue.

### INTRODUCTION

Caffeine is probably the most frequently ingested pharmacologically active substance in the world. It is found in common beverages (coffee, tea, soft drinks), in products containing cocoa or chocolate, and in medications. Caffeine an alkaloid of the methylxanthine family is a naturally occurring substance found in the leaves, seeds or fruits of over 63 plants species worldwide [1]. In its pure state, it is an intensely bitter white powder. Its chemical formula is C<sub>8</sub>H<sub>10</sub>N<sub>4</sub>O<sub>2</sub>, its systematic name is 1,3,5-trimethylxanthine [2]. Caffeine is a pharmacologically active substance and depending on the dose, can be a mild central nervous system stimulant. Caffeine does not accumulate in the body over the course of time and is normally excreted within several hours of Consumption [3]. Other naturally occurring

methylxanthines include theobromine and theophylline. Methylation of theobromine forms caffeine. [1].

Caffeine is used as an ergogenic aid because multiple well-controlled experiments have found that moderate doses of caffeine (3-6 mg/kg) can improve performance in athletes [4,5]. Its use by athletes is actually favoured since it has been removed on the 1st of January 2004 by the World Anti-Doping Agency (WADA) from its list of banned substances [6]. The mechanism for the caffeine-improved performance is not clear but several possibilities have been proposed such as the antagonism of adenosine receptors [7], the attenuation of effort perception or reduction of muscle pain [8,9] and the increase in catecholamine release [10-13].

There have been numerous reports that caffeine is an ergogenic aid; ingestion of the drug has been shown to increase endurance, particularly in prolonged exercise



lasting 30-120 min [14-16]. Coffee is certainly the most common mode of caffeine ingestion, and it also can provide caffeine in a more concentrated form than do other foodstuffs [17,18]. Caffeine and coffee ingestion have been shown to result in similar hemodynamic responses in resting participants [19]. The aim of the present study was to study the effects of caffeine in coffee on physical performance, exercise intensity and muscle strength in young healthy adults.

## MATERIALS AND METHODS

Participants in this case control study were twenty young healthy males aged between 17-24 yrs studying in 1st MBBS. They were non athletes, non regular users of caffeine, non smokers & apparently healthy. Males with history of smoking, asthma, hypertension, congenital heart disease, allergy, diabetes were excluded from the study. The study was carried out in the Exercise Physiology Unit of Department of Physiology, Jawaharlal Nehru Medical College, Sawangi (Meghe), Wardha, Maharashtra in the month of January. The participants were informed both verbally and in writing about the nature of the experiments and the protocol was approved by the Ethics Committee of the Institution.

### Experimental protocol

Each participant reported to the Exercise unit on three occasions. On the first occasion they performed an incremental  $\dot{V}O_2$  max test on a treadmill (RMS Polyrite, Chandigarh). Subsequently, on a separate day, the participants performed a practice trial consisting of 20-30 min running on the treadmill at a workload predicted to require 80% of the  $\dot{V}O_2$  max. On the same day participants basic parameters such as age, weight, height, body mass index and heart rate maximum (HRmax) were recorded as given in table (1). Maximal voluntary isometric arm flexion strength was measured by handgrip dynamometer and the best result of three readings was recorded. Rate of perceived exertion by Borg's 10 point scale [20] was recorded at 3 min intervals throughout the exercise and exhaustion time was recorded on treadmill. Time to exhaustion was determined on a treadmill (the time that the subject could no longer maintain exercise intensity or reached volitional exhaustion).

The participants were instructed to refrain from consuming caffeine for 48 h prior to each testing session. A list of food and beverages containing caffeine was provided to each participant in order to inform them of what products they should avoid. Participants were also asked to keep a diary documenting

their diet 24 h prior to the first testing session and to replicate this diet before next visit.

The participants arrived at the laboratory on all the occasions at the same time (7:30 or 8:15 AM). When the participants reported to the Exercise unit after seven days they were given filter coffee with desired amount of caffeine needed (5mg/kg) dissolved in 200ml warm water and added artificial sweetener (sugar free gold i.e aspartame). The dose of 5.0 mg·kg<sup>-1</sup> was selected based on previous studies that used this dose and reported performance benefits [21].

Participants relaxed for 60 min, followed by an active warm up they were instructed to run on the treadmill at a speed and slope calculated to require 80% of  $\dot{V}O_{2max}$  until voluntary exhaustion. Endurance i.e exhaustion time, Rate of perceived exertion, handgrip strength test and HRmax was recorded as done previously.

### Estimation of $\dot{V}O_2$ max

$\dot{V}O_2$  max was calculated by following equation,  $\dot{V}O_2$  max = resting component [1MET {3.5 ml O<sub>2</sub>/kg/min} + Horizontal component {speed (m/min) x oxygen consumption of vertical movement}] (40).

### Statistical Analysis

All results are expressed as means  $\pm$  SD. Students Paired 't' test was applied and the level of significance was set at  $P < 0.05$ .

## RESULTS

All results refer to 15 subjects, unless otherwise indicated. Statistical analysis pertains to results obtained in 15 subjects. There was a significant difference in all the parameters before and after caffeine ingestion. All the results are shown in table 2.

### Exhaustion time and $\dot{V}O_2$ max

Ingestion of caffeine significantly increased the time to exhaustion in all subjects (545.10 sec  $\pm$  3.50 SD to 608  $\pm$  42.01 SD) with  $p < 0.05$ . There was also a significant difference in  $\dot{V}O_2$  max after caffeine ingestion (35.36  $\pm$  4.01 SD to 40.33  $\pm$  3.70,  $p < 0.05$ ).

### Handgrip and RPE scale

There was a significant difference in the isometric muscle strength by handgrip dynamometer and in the rate of perceived exertion before and after caffeine ingestion (86.25  $\pm$  3.50 SD to 88.25  $\pm$  3.11 SD with  $p < .05$  and 4.45  $\pm$  0.51 to 6.85  $\pm$  0.87,  $p < .05$ ) respectively.

There was also a significant difference in the HR max after caffeine ingestion (176.50  $\pm$  6.01 to 179.40  $\pm$  5.95) with  $p < 0.05$ .

**Table 1. Descriptive statistics for age, height, weight and BMI**

Parameters	Mean	SD	Minimum	Maximum
Age(yrs)	18.65	0.74	18.00	20.00
Height(cm)	70.15	3.09	65.00	75.00
Weight(kg)	173.95	4.58	166.00	182.00
BMI(kg/m <sup>2</sup> )	23.25	0.67	21.80	24.20



All values are in mean-SD



**Table 2. Comparison of data before and after ingestion of coffee**

	Mean		N		Std. Error Mean	
	Before	After	Before	After	Before	After
Handgrip	86.25±3.50	88.25±3.118*	20	20	0.78	0.69
RPE scale	4.45±0.51	6.85±3.70*	20	20	0.11	0.19
VO2 max	35.36±4.01	40.33±3.70*	20	20	0.89	0.82
Hrmax	176.50±6.01	179.40±5.95*	20	20	1.34	1.33
Exhaustion time	545.10±40.58	608.85±42.01*	20	20	9.07	9.39

All the values are in mean ± SD,\*indicates a significant difference with p<0.05.



**DISCUSSION**

Research on the physiological effects of caffeine as it relates to athletic performance is extensive. Caffeine research in specific areas of interest, such as endurance, strength, team sport, recovery, and hydration is vast and at times, conflicting [22]. Therefore the present study was undertaken which included 20 young healthy adult males studying in first MBBS, to study the effects of caffeine in coffee on exhaustion time, VO2 uptake, muscle strength, rate of perceived exertion and heart rate maximum. Overall, results from this study showed that caffeine in coffee resulted in significant increases in all the parameters examined.

Borg demonstrated that the general perception of physical exertion comes from the integration of different symptoms arising from active muscles, cardiovascular and respiratory systems, joints, perspiration, possible pain, dizziness etc. The Borg CR-10 scale is a category scale with ratio properties consisting of numbers related to verbal expressions, which allows rate comparison between intensities as well as a determination of intensity levels, it has been used for more than two decades. The use of this scale is relatively simple and cost effective. In our study we used this scale which demonstrated that the rate of perception of physical exertion increased after caffeine ingestion. There are studies that have failed to detect statistically significant changes in the perceptual response to exercise following caffeine ingestion, for example, both Tarnopolsky et al [23] and Trice et al [24] found that

caffeine ingestion reduced RPE by approximately 16% and 6% in comparison to placebo when six and eight subjects ran and cycled between 70% and 90% VO2max, respectively. A meta-analytic study done by M. Doherty & PM. Smith quantifies suggestions in the literature that caffeine has a noticeable effect on RPE, which is in support of findings in our study.

Caffeine ingestion resulted in augmented exercise performance as indicated by a significantly higher time to exhaustion and maximal O2 consumption as seen in our study. These results are consistent with those reported by other authors.

Multiple mechanisms have been proposed to explain the effects of caffeine supplementation on sport performance. However, several extensive reviews have stated that the most significant mechanism is that caffeine acts to compete with adenosine at its receptor sites. Another possible mechanism through which caffeine may improve endurance performance is by increasing the secretion of β-endorphins. Laurent et al. [25] demonstrated that when compared to the placebo group caffeine consumption (6 mg/kg) significantly increased plasma β-endorphin concentrations following two hours of cycling at 65% VO2peak and a subsequent bout of high intensity sprint activity. It has been established that plasma endorphin concentrations are enhanced during exercise and their analgesic properties may lead to a decrease in pain perception [26].



Research has also demonstrated that caffeine may result in alterations of neuromuscular function and/or skeletal muscular contraction [27,28]. Kalmar and Cafarelli [27] indicated a moderate dose of caffeine (6 mg/kg) significantly enhanced both isometric leg extension strength as well as the time to fatigue during a submaximal isometric leg extension.

Whatever caffeine's mechanism of action, one consistent outcome of caffeine ingestion during exercise testing, regardless of mode, intensity, or duration of exercise, is an alteration in participants' perceptual response. This alteration has been manifest as either an increase in work output at a given ratings of perceived exertion (RPE) or effort sense [29-31] or, more typically, a reduced RPE at a constant exercise [32,-35].

Various methods of caffeine supplementation have been explored and results have provided considerable insight into appropriate form and dosage of the compound. One of the most acknowledged studies, published by Graham et al. [21] demonstrated a range of effects when caffeine (at 4.45 mg/kg) was consumed in varying forms. In their study, aerobically conditioned runners performed five treadmill runs to exhaustion at approximately 85%  $VO_{2max}$  after receiving one of the following treatments 60 minutes prior: caffeine capsules plus water, regular coffee, decaffeinated coffee, decaffeinated coffee plus caffeine in capsule form, and placebo. Caffeine in capsule form significantly increased work capacity allowing them to run an additional 2-3 km [21], as compared to the four other treatments. Caffeine in capsule form significantly increased work capacity allowing them to run an additional 2-3 km [21], as compared to the four other treatments.

As such, McLellan and Bell [36] examined whether a morning cup of coffee just prior to anhydrous caffeine supplementation would have any negative impact on the compound's ergogenic effect. Subjects were physically active and considered to be moderate-to-high daily consumers of caffeine. The results of their study indicated that caffeine supplementation significantly increased exercise time to exhaustion regardless of whether caffeine in anhydrous form was consumed after a cup of regular or decaffeinated coffee [36].

## REFERENCES

1. Wanyika HN, Gatebe EG, Gitu LM, Ngumba EK and Maritim CW. (2010). Determination of caffeine content of tea and instant coffee brands found in the Kenyan market. *African Journal of Food Science* 4(6), 353-358.
2. Aurnaud MJ. (1987). The pharmacology of caffeine. *Prog Drug*, 31, 273.
3. Barone JJ, Roberts HR. (1996). Caffeine consumption. *Food Chem Toxicol*, 34, 119.
4. Graham T. (2001). Caffeine and exercise, metabolism, endurance and performance. *Sports Medicine*, 31, 785-807.
5. Flinn S, Gregory J, Mcnaughton L, Tristram S and Davies P. (1990). Caffeine ingestion prior to incremental cycling to exhaustion in recreational cyclists. *International Journal of Sports Medicine*, 11, 188-193.
6. World Anti-Doping Agency. (2008). *The world anti-doping code. The 2009 prohibited list. International standard*. Retrieved October 24, 2008, from URL, [http://www.wada-ama.org/rtecontent/document/2009\\_Prohibited\\_List\\_ENG\\_Final\\_20\\_Sept\\_08.pdf](http://www.wada-ama.org/rtecontent/document/2009_Prohibited_List_ENG_Final_20_Sept_08.pdf)
7. Davis J, Zhao Z, Stock H, Mehl K, Buggy J and Hand G. (2003). Central nervous system effects of caffeine and adenosine on fatigue. *American Journal of Physiology. Regulatory, Integrative and Comparative Physiology*, 284, R399-404.

McNaughton et al. [37] reported the effects of a moderate dose of caffeine (6 mg/kg) on 1-hour time trial performance. This investigation was unique to the research because, while continuous, the protocol also included a number of hill simulations to best represent the maximal work undertaken by a cyclist during daily training. The caffeine condition resulted in the cyclists riding significantly further during the hour-long time trial, as compared to placebo and control. The time trial performance was improved 4-5% by the caffeine treatment over the other two treatments [37]. Demura et al. [38] examined the effect of coffee, which contained a moderate dose of caffeine at 6 mg/kg, on submaximal cycling. Subjects consumed either caffeinated or decaffeinated coffee 60 min prior to exercise. The only significant finding was a decreased RPE for the caffeinated coffee as compared to the decaffeinated treatment [38].

Coffee contains multiple biologically active compounds; however, it is unknown if these compounds are of benefit to human performance [39-45]. However, it is apparent that consuming an anhydrous form of caffeine, as compared to coffee, prior to athletic competition would be more advantageous for enhancing sport performance. Nevertheless, the form of supplementation is not the only factor to consider as appropriate dosage is also a necessary variable.

Reviews of the physiological effects of caffeine commonly give an approximate equivalent of various food items such as coffee, tea, colas, etc. The implication would appear to be that ingesting the same amount of caffeine via a food source is as effective as ingesting pure caffeine.

## CONCLUSION

The results of this study support the use of coffee with an adequate dose of caffeine (5mg/kg BW) before exercise can help to improve physical performance in physically active adults. Competing with adenosine to its receptor site, secretion of  $\beta$ -endorphins or alterations of neuromuscular function and/or skeletal muscular contraction could be the several possible mechanisms that may account for this improvement. Further research is required to support the same.



8. Doherty M and Smith P. (2004). Effects of caffeine ingestion on exercise testing, a meta-analysis. *International Journal of Sport Nutrition and Exercise Metabolism*, 14, 626-646.
9. O'connor P, Motl R, Broglio S and Ely M. (2004). Dose-dependent effect of caffeine on reducing leg muscle pain during cycling exercise is unrelated to systolic blood pressure. *Pain*, 109, 291-298.
10. Greer F, Friars D and Graham T. (2000). Comparison of caffeine and theophylline ingestion, exercise metabolism and endurance. *Journal of Applied Physiology*, 89, 1837-1844.
11. Graham T, Helge J, Maclean D, Kiens B and Richter E. (2000). Caffeine ingestion does not alter carbohydrate or fat metabolism in human skeletal muscle during exercise. *The Journal of Physiology*, 529, 837-847.
12. Jackman M, Wendling P, Friars D and Graham T. (1996). Metabolic catecholamine and endurance responses to caffeine during intense exercise. *Journal of Applied Physiology*, 81, 1658-1663.
13. Van Soeren MH and Graham TE. (1998). Effect of caffeine on metabolism, exercise endurance and catecholamine responses after withdrawal. *J Appl Physiol*, 85,1493-1501.
14. Reilly T, Orme M, Graham TE. (1997). The possible actions of methylxanthines on various tissues in. *The Clinical Pharmacology of Sport and Exercise*, eds Reilly T., Orme M. (Elsevier Science, Amsterdam), 257-270.
15. Graham TE, Rush JWE, VanSoeren MH. (1994). Caffeine and exercise, metabolism and performance. *Can J Appl Physiol*. 19, 111-138.
16. Graham TE, Spriet LL. (1991). Performance and metabolic responses to a high caffeine dose during prolonged exercise. *J Appl Physiol*, 71, 2292-2298.
17. Garattini S, Arnaud MJ. (1993). Metabolism of caffeine and other components of coffee. in *Caffeine, Coffee, and Health*, ed Garattini S. (Raven, New York), 43-96.
18. Lamb DR, Williams MH, Conlee RK. (1991). Amphetamine, caffeine, and cocaine. In *Ergogenics—Enhancement of Performance In Exercise and Sport*, eds Lamb D. R., Williams M. H. (Brown, Ann Arbor, MI), 285-330.
19. Casiglia E, Bongiovi S, et al. (1991). Haemodynamic effects of coffee and caffeine in normal volunteers, a placebo-controlled clinical study. *J Intern Med*, 229, 501-504.
20. Borg GAV. (1982). Psychophysical bases of perceived exertion. *Medicine and Science in Sports and Exercise*, 14, 377-81
21. Graham TE, Hibbert E, Sathasivam P. (1998). Metabolic and exercise endurance effects of coffee and caffeine ingestion. *J Appl Physiol*, 85, 883-889.
22. Fredholm BB, Battig K, Holmen J, Nehlig A, Zvartau EE. (1999). Actions of caffeine in the brain with special reference to factors that contribute to its widespread use. *Pharmacol Rev*, 51, 83-133.
23. Tarnopolsky MA, Atkinson SA, Macdougall JD, Sale DG, Sutton JR. (1989). Physiological responses to caffeine during endurance running in habitual caffeine users. *Med Sci Sports Exerc*, 21, 418-424.
24. Trice I, Haymes EM 1995. Effects of caffeine ingestion on exercise-induced changes during high-intensity, intermittent exercise. *Int J Sports Nutr*, 5, 37-44.
25. Sokmen B, Armstrong LE, Kraemer WJ, Casa DJ, Dias JC, Judelson DA, Maresh CM. (2008). Caffeine use in sports, Considerations for the athlete. *J Strength Cond Res*, 22, 978-986.
26. Spriet LL, Gibala MJ. (2004). Nutritional strategies to influence adaptations to training. *J Sports Sci*, 22, 127-41.
27. Laurent D, Schneider KE et al. (2000). Effects of caffeine on muscle glycogen utilization and the neuroendocrine axis during exercise. *J Clin Endocrinol Metab*, 85, 2170-75.
28. Grossman A, Sutton JR. (1985). Endorphins, What are they? How are they measured? What is their role in exercise? *Med Sci Sports Exerc*, 17,74-81.
29. Kalmar JM, Cafarelli E. (1999) Effects of caffeine on neuromuscular function. *J Appl Physiol* 87, 801-808.
30. Lopes JM, Aubier M, Jardim J, Aranda JV, Macklem PT. (1983). Effect of caffeine on skeletal muscle function before and after fatigue. *J Appl Physiol Respirat Environ Exercise Physiol*, 54, 1303-1305.
31. Ivy JL, Costill DL, Fink WJ, Lower RW. (1979). Influence of caffeine and carbohydrate feedings on endurance performance. *Med Sci Sports*, 11, 6-11
32. Cole KJ, Costill DL, Starling RD, Goodpastor BH, Trappe SW, Fink WJ. (1996). Effect of caffeine ingestion on perception of effort and subsequent work production. *Int J Sports Nutr*, 6, 14-23.
33. Plaskett CJ, Cafarelli E. (2001). Caffeine increases endurance and attenuates force sensation during submaximal isometric contractions. *J Appl Physiol*, 91, 1535-1544.
34. Costill DL, Dalsky GP, Fink WJ. (1978). Effects of caffeine on metabolism and exercise performance. *Med Sci Sports*, 10, 155-158.
35. Giles D, Maclaren D. (1984). Effects of caffeine and glucose ingestion on metabolic and respiratory functions during prolonged exercise. *J Sports Sci*, 2, 35-46.
36. Casal DC, Leon AS. (1985). Failure of caffeine to affect substrate utilization during prolonged running. *Med Sci Sports Exerc*, 17, 174-179.
37. MacIntosh BR, Wright BM. (1995). Caffeine ingestion and performance of a 1500m swim. *Can J Appl Physiol*, 20, 168-177.



38. McLellan TM, Bell DG. (2004). The impact of prior coffee consumption on the subsequent ergogenic effect of anhydrous caffeine. *Int J of Sport Nutr Exerc Meta*, 14,698-708.
39. McNaughton LR, Lovell RJ, Siegler JC, Midgley AW, Sandstrom M, Bentley DJ.(2008) The effects of caffeine ingestion on time trial cycling performance. *J Sports Med Phys Fitness*, 48, 320-5.
40. Demura S, Yamada T, Terasawa N. (2007). Effect of coffee ingestion on physiological responses and ratings of perceived exertion during submaximal endurance exercise. *Perceptual Motor Skills*, 105, 1109-16.
41. Natella F, Nardini M, Giannetti I, et al. (2002). Coffee drinking influences plasma antioxidant capacity in humans. *J Agric Food Chem*, 50, 6211- 6.
42. William D, Mcardle FI, Katch V, Katch I. (2000). Energy expenditure during walking, jogging, running & swimming. Chapter 10, 213,7<sup>th</sup> edition].
43. Erica R Goldstein et al. (2010). International society of sports nutrition position stand, caffeine and performance. *Journal of the International Society of Sports Nutrition*, 7, 5.
44. Guillermo J Olcina et al. (2006). Effect of caffeine on oxidative stress during maximum incremental exercise. *Journal of Sports Science and Medicine*, 5, 621-628.
45. Doherty M, Smith PM. (2005). Effects of caffeine ingestion on rating of perceived exertion during and after exercise, a meta-analysis. *Scand J Med Sci Sports*, 15, 69–78.

