




AMERICAN JOURNAL OF BIOLOGICAL AND PHARMACEUTICAL RESEARCH



Journal homepage: www.mcmed.us/journal/ajbpr

GUT MICRO-FLORA OF EARTHWORMS: A REVIEW

B.Govindarajan^{1*} and V.Prabakaran²

¹Department of Zoology, VHNSN College (Autonomous), Virudhunagar, Tamil Nadu, India.

²Department of Zoology, Government Arts College, Karur, Tamil Nadu, India.

Article Info

Received 29/07/2014

Revised 16/08/2014

Accepted 19/08/2014

Key words: -

Earthworm, Gut,
Bacteria, Fungi and
Microflora.

ABSTRACT

This review paper describes the different species of earthworm gut microflora. Laboratory cultures and experiments were carried out to investigate the fate of bacteria during and after passage through the intestinal tract of detritivore earthworms. The gut of many earthworm species contains a complex and mutualistic microbial community that usually assist the earthworms with digestion and vermicompost. Finally this review discussed the earthworm gut microorganisms.

INTRODUCTION

In the 21th century, also why we concentrate on earthworms the answer is today earthworms widely accepted as ecosystem engineer. Earthworms increase microbial activities by providing in their gut mucus consisting of energetic and easily metabolizable compounds [1] and considerable physico-chemical conditions: neutral pH, high moisture and ideal temperature conditions [2]. Earthworms seem to have poor proper enzymatic systems and they appear to rely upon the ingested soil microorganisms to degrade soil organic matter.

A major part of earthworm activity on soil properties is contributed to interactions with soil vicinity microorganisms [3]. However, these interactions are still not clearly understood, including the effect of gut passage on the community structure of ingested soil microorganisms. Interactions between earthworms and soil microorganisms are key character for soil processes such as decomposition and transformation of plant residue, humus formation and the formation of the pool of nutrient

elements and microbial communities. The wide spectrum of these interactions makes it possible to speak about a close relationship between earthworms and soil microorganisms. Soil fungi, bacteria etc are assumed to be the main source of food for earthworms [4]. While feeding, earthworms regulate the growth of soil microorganisms by eating some microbial populations and providing ideal conditions for the growth of others in their digestive tract and in casts [5]. The gut of many soil organisms (microinvertebrates and macroinvertebrates) contains microbial communities that usually helpful in the digestion. These microbial-animal relationships create mutualisms. Earthworms are also having a mutualistic relationship with soil microorganisms (Bacteria, Fungi, etc) passing through their digestive tract, but the nature and role of the microbiota inhabiting their gut are virtually unknown [6]. However, little is understood on the interactions between earthworm and soil microorganisms, including the gut microbial community. The goal of the present review work was to list out the gut microflora of different species of earthworms.

Corresponding Author

B.Govindarajan

Email:- bgmphilbed@gmail.com

Why Earthworm

Earthworms are worldwide considered as an excellent bioindicators of the relative health of soil vicinity and possess a number of qualities that predispose them for



use in monitoring terrestrial ecosystems [7]. Earthworms are decomposers, aerators, crushers, mixers, chemically degraders and biologically stimulators in the soil vicinity. They effectively harness the beneficial soil microflora, destroy soil pathogens and convert organic wastes (Domestic waste, Industrial waste, Hospital waste, etc) into vitamins, enzymes, antibiotics, growth hormones and protein rich casts.

Earthworm Gut

Earthworm guts considered as ideal habitats for microorganisms (bacteria, etc), because several studies showed increased microbial numbers in the guts versus the soil, in which earthworms were living [8]. Some previous studies proposed that the earthworm gut microbial community is qualitatively not much different from the microbial community in the surrounding soil vicinity [9], later studies found significant differences for selected phylogenetic groups or functional guilds of microorganisms, eg: Proteobacteria [10], Actinobacteria, denitrifiers or cellobiose utilizers [11,12].

Earthworms gut performs a unique vicinity subsystem of soil vicinity. The earthworm gut has stable conditions different from surrounding vicinity. Earthworm gut is a straight tube bioreactor, which maintains a stable temperature through novel temperature regulatory mechanisms, thus accelerating the rates of the bioprocesses and preventing enzyme inactivation caused by high temperatures. Earthworm's gizzard is a novel colloidal mill in which the feed is ground into particles smaller than 2 mm giving, thereby, enhancing surface area for microbial processing [13].

Earthworm gut is basically an effective tubular structure extending from mouth to the anus; its different regions are the muscular pharynx, oesophagus, intestine and associated digestive glands. The gut contents usually comprise mucus, organic and mineral matter. An analysis of gut contents in earthworm revealed the occurrence of different kinds of symbiont like microfungi, bacteria, protozoa, etc; most microfungi species are present in the foregut, gradually decreased in number in the mid and hindgut with fewest in freshly laid casts [14]. It is well established that the earthworm gut provides ideal conditions for the development of microorganisms (bacterial & fungi colonies) since earthworm casts contain significantly higher counts of bacteria than in the surrounding soil [15-19].

The rate of food transit along the digestive tract of earthworms is very high. Therefore, it appears that earthworms quickly kill and digest the soil microorganisms consumed. It has previously been shown that the digestion of microorganisms in the gut of invertebrates begins with the quick death of certain microbial cells caused by specific killing agents of nonprotein nature [20].

Microorganisms may constitute a very important part of the diet of earthworms, which can feed on them selectively [21,22]. The gut environment is anoxic, pH 6.9 with about 50% water content. The gut bacteria are enriched in total carbon, organic carbon and total nitrogen with a carbon to nitrogen ratio of 7 [23]. The bacterial counts in guts/vermicompost were higher than the surrounding soil ecosystem [24, 25] and as the organic matter ingested passes through the gut, it undergoes biochemical changes effected by gut-inhibiting bacteria. Therefore there is a greater role played by the DNA of microorganisms and earthworm species. The genetic makeup of the strains and environment has a profound influence on the efficiency of earthworms in the bioconversion process [26] and in the assessment of diversity and community composition [27].

The earthworm gut could be used as an environment and nutrition by microorganisms and gut derived enzymes could affect the microbes on the soils particles [28]. On the other hand, the earthworm is migrating in the soil and thus soil flows through earthworm gut. In the gut of the earthworms (*Eisenia fetida*) the microbial composition was changed towards the increasing numbers of non-spore forming bacteria and decreasing numbers of spore forming bacteria, this also resulted in enhanced levels of nitrogen fixation [29]. A number of novel N₂O producing species of bacteria from genera dechloromonas (Bata Proteo Bacteria), from genera dechloromonas (Cytophaga Flara bacteria group Bacteroidetes).

Earthworm activity does not only mediate macroaggregate formation, but also microaggregate formation [30]. Based on thin sections of the earthworm gut, casts and control soil from earthworm microcosms, several studies have shown that during gut transit organic materials are intimately mixed and become encrusted with the mucus to create new nuclei for microaggregate formation [31].

The digestive system of earthworm consists of a pharynx, oesophagus and gizzard (reception zone) followed by an anterior intestine that secretes enzymes and a posterior intestine that absorbs nutrients. Earthworm activity does not only mediate macroaggregate formation, but also microaggregate formation. Based on thin sections of the earthworm gut, casts and control soil from earthworm microcosms, several studies have shown that during gut transit organic materials are intimately mixed and become encrusted with the mucus to create new nuclei for microaggregate formation.

Earthworm Gut Bacteria

The diversity of 8 bacterial groups from fresh soil and gut of the earthworms *L. terrestris* and *Aporrectodea caliginosa* were studied by single strand conformation polymorphism (SSCP) analysis using both newly designed



16S rRNA gene specific primer sets targeting Alphaproteo bacteria, Betaproteo bacteria, Gammaproteo bacteria, Deltaproteo bacteria, Bacteroides, Verrucomicrobia, Planctomycetes and Firmicutes as a conventional universal primer set for SSCP, with RNA and DNA as templates. Whereas using fluorescence in situ hybridization Bacteroidates, Alphaproteo bacteria and Betaproteo bacteria were predominant in communities from the soil and worm cast samples, some specific bacterial taxonomic groups maintain their diversity and even increase their relative numbers during transit through the gastrointestinal tract of earthworms [32].

Lumbricus terrestris, *L. friend*, *Aporrectodea caliginosa* and *A. longa* contain ecological group specific gut wall associated bacterial communities. The abundance of specific gut wall associated bacteria, including proteobacteria, firmicutes and an actinobacterium was dependent on the ecological group. Bacteria are of minor importance in the diet, algae are of moderate importance; protozoa and fungi are major sources of nutrients. Worms, produced under sterile conditions, could live on individual cultures of certain bacteria, fungi and protozoa, but grew best on various mixtures of microorganisms. Symbiotic interactions between earthworms and microorganisms break down and fragment organic matter progressively, finally incorporating it into water stable aggregates.

Valle Molinares *et al* 2007 identified 7 species of bacteria from the genus Bacillus (*B. insolitus*, *B. megaterium*, *B. breris*, *B. pasteurii*, *B. sphaericus*, *B. thuringiensis* and *B. pabuli*) within the intestine of *Onychochaeta borincana*. All these species are typical soil bacteria. In addition, it was found that the microbial weight of the intestinal region decreased from the anterior to posterior section. For studies on bacteria within the intestine of earthworms, diverse methods and techniques have been used which have helped in identifying species of the genus Bacillus, Pseudomonas, Klebsiella, Azotobacter, Serratia, Aeromonas and Enterobacter [33,34].

Earthworm gut environment may act as specific filter as well as a fermenter for some soil bacteria and fungi with those bacterial cells that survive passage through the midgut can even multiply in the hindgut. Spores of some fungi that survived in the midgut vicinity of earthworms.

Researchers have attempted to study earthworm gut microbes using direct culture methods [eg: 35,36] and electron microscopy [37]. Microbial numbers in the earthworm gut are higher than those in surrounding soil and denitrification, but not methane emission, occurs in the gut of *Aporrectodea caliginosa* and *Lumbricus rubellus* [38]. In addition, gene clone libraries of bacteria tightly associated with the gut wall were different from those in the gut content [39]. These findings suggest that the earthworm gut is favourable for the growth and activity of certain bacterial species.

36 reported the first Scanning Electron Microscopy (SEM) study on the digestive tube of earthworms. They find the coccid bodies from the foregut of *O. cyaneum*. In *O. borincana* they were only seen in the hindgut. Earthworm revealed the presence of cocci, rod-shaped bacteria and filamentous microorganisms in *Lumbricus terrestris* L., 1758 and *Octolasion cyaneum*.

The key role of the microorganisms in the gut of earthworms is not clearly understood: indigenous bacteria are not generally found in earthworms while, if present they may aid digestive processes. Seven different species of *Bacillus* have been identified from the gut of *O. Borincana* [40]. Apparently, the presence of these microorganisms in the intestinal tube may be a result of soil ingestion and do not seem to assist in its digestion. Although rod-shaped bacteria were observed associated to the intestinal wall of *O. borincana*.

The profiles of bacteria are very closely similar in soil and in the gut contents of many earthworm species, suggesting an absence in the gut of any indigenous bacterial groups. In the time of scanning of the gut surface of these two species (*Lumbricus terrestris* and *Octolasion cyaneum*), only two rod shaped organisms were found. In contrast, the Transmission Electron Microscopy (TEM) investigation of *L. terrestris* hindgut showed several rod-shaped bacteria over a relatively small area.

Earthworms are ubiquitous soil invertebrates that ingest large amounts of mineral soil and organic matter containing a variety of microorganisms [41]. [42] concluded that a field population of 75gm⁻² per ha. The effects on bacteria of the passage through the earthworm intestinal tract may have a major impact on the composition of the soil bacterial community and so gut passage should be taken into consideration when assessing the risk of releasing non-indigenous, eg genetically engineered, bacteria into terrestrial ecosystems.

Denitrification in the earthworm gut is involved in the in vivo emission of N₂O by earthworms, cultured denitrifiers occur in high numbers in the earthworm gut [43-45]. Earthworms can directly regulate microbial population (Bacteria, Fungi) by consuming large amount of soil. This leads to elimination of some microorganisms and proliferation of others in the digestive tract of earthworm [46].

Microorganisms are a vital food component of soil invertebrates including earthworms. The importance to fungi and bacteria as important sources of food is indicated by several phenomena. Soil animals avoid consuming fresh leaf litter, which could be toxic, the animals themselves have no intrinsic capacity to digest cellulose and they depend on microorganisms as sources of essential amino acids [47,48].

It was proposed that earthworms derive more of its energy and nutrients from gut specific microbiota than from microbiota already present in the ingested soil [49]. [50]



Who found in a series of tests that the total number of microorganisms (bacteria, fungi) in a alimentary tract or in the fresh excrements of earthworms were from half to 6 times higher in comparison with the surrounding soil.

Fungal flora

Microfungi in the casts of earthworms, *Perionyx millardi*, *Eudrilus eugeniae*, *Lampito mauritii* and certain other earthworm species are available. But, the scientific knowledge available on the fungal flora in the casts of the earthworm, *Perionyx ceylanensis* Mich. *Aspergillus fumigates*, *Mucor circinelloides*, *F. circinelloides* and *Penicillium expansum* were dominant in the intestine of earthworms [51].

Earthworms and Microbes interaction in Vermicompost formation

Earthworms promote the growth of beneficial decomposer microbes (bacteria, actinomycetes and fungi) in waste biomass [52]. They hosts millions of decomposer microbes in their gut which is described as little bacterial

factory. They devour on microbes and excrete them out (many time more in number than they ingest) in soil along with nutrients nitrogen (N) and phosphorus (P) in their excreta. The nutrients N and P are further used by the microbes for multiplication and vigorous action.

Earthworms and microbes act symbiotically and synergistically to accelerate and enhance the decomposition of the organic matter in the waste. It is the microorganisms that break down the cellulose in the food waste, grass clippings and the leaves from garden wastes [53].

CONCLUSION

In conclusion, we would like to conclude that the earthworm gut is an ideal vicinity for microorganisms such as bacteria, fungi etc.

ACKNOWLEDGMENT

This work was supported by the VHNSN College Management and we wish to express gratitude lord Shiva for priceless assistance.

REFERENCES

1. Martin A, Cortez J, Barosis I and Lavelle P. Les mucus intestinaux de Var de Terre, moteur de leurs interactions avec la microflore. *Rev Ecol Biol Sol*, 24, 1987, 549-558.
2. Barois I and Lavelle P. Changes in respiration rate and some physico-chemical properties of a tropical soil during transit through *Pontoscolex corethrurus* (Glossoscolecidae, Oligochaeta). *Soil Biol Biochem*, 18, 1986, 539-541.
3. Edwards CA and Fletcher KE. Interactions between earthworms and microorganisms in organic matter breakdown. *Agric Ecosyst Environ*, 24, 1988, 235-247.
4. Byzov BA. Zoomicrobial interactions in soil. Extended Abstract of Doctoral (Biol.) Dissertation, Moscow, Izd-vo "Maks Press", 2003.
5. Tiunov AV and Scheu S. Microfungai, communities in soil, litter and cast *Lumbricus terrestris* L. (Lumbricidae), a Laboratory experiment. *Appl Soil Ecol*, 14, 2000, 17-26.
6. Lavelle P and Spain AV. Soil ecology. Kluwer Academic Publishers, Dordrecht. The Netherlands. 2001, 684.
7. Vandecasteele B, Samyn J, Quataert P, Muys B, Tack FMG. Earthworm biomass as additional information for risk assessment of heavy metal biomagnification, a case study for dredged sediment-derived soils and polluted floodplain. *Soils Environ Pollut*, 129, 2004, 363-375.
8. Wolter C and Scheu S. Changes in bacterial numbers and hyphal lengths during the gut passage through *Lumbricus terrestris* (Lumbricidae, Oligochaeta). *Pedobiologia*, 43, 1999, 891-900.
9. Parle JN. Microorganisms in the intensions of earthworms. *J Gen Microbiol*, 31, 1963, 1-11.
10. Schonholzer F, Hahn D, Zarda B and Zeyer J. Automated image analysis and in situ hybridization as tools to study bacterial populations in food resources, gut and cast of *Lumbricus terrestris* L. *J Microbiol Methods*, 48, 2002, 53-68.
11. Kritufek V, Ravasz K and Pizl V. Actinomycete communities in earthworm guts and surrounding soil. *Pedobiologia*. 37, 1993, 379-384.
12. Ihssen J, Horn M A, Matthies C, Gossner A, Schramm A and Drake HL. N₂O producing microorganisms in the gut of the earthworm *Aporrectodea caliginosa* are indicative of ingested soil bacteria. *Appl. Environ. Microbiol.* 69, 2003, 1655-1661.
13. Karthikeyan S, Ramesh P T, Nagamani B and Kumar GM. Earthworms. *Agro, India*. 7, 2004, 34-35.
14. Dash MC, Senapati BK and Mishra CC Nematode feeding by tropical earthworms. *Oikos*, 34, 1980, 322-325.
15. Edwards CA and Lofty Biology of earthworms. Chapman and Hall, London, 1977, 129-147.
16. Kale RD, Mallesh BC, Kubra B and Bhagyaraj DJ. Influence of vermicompost application on available micronutrients and selected microbial populations in paddy field. *Soil Biol & Biochem*, 24, 1992, 1317-1320.
17. Munnoli PM. A study on management of organic solid waste of agro based industries through vermiculture biotechnology. ME Thesis, TIET Patiala, India, 1998, 11-30.



18. Bhattacharyya KK, Mukhopadhyay N, Mukharjee D, Das SK. Comparative efficiency of improved compost techniques. In, Jana BB, Banerjee RD, Guterstam B, Heeb J (Eds) Proceedings of International Conference Waste Recycling & Resource Management in Developing World, Sapana Printing Works, Kolkatta, India, 2000, 219-224.
19. Munnoli PM. Management of industrial organic solid wastes through vermiculture biotechnology with special reference to microorganisms. PhD, Thesis, Goa University, India, 2007, 1-334.
20. Byzov BA, Vu Nguen Thanh, Bab Eva IP, Tretyakova EB, Dyvak IA and Rabinovich Ya M. Killing and Hydrolytic activities of the gut fluid of the Millipede *Pachyiulus flavipes* CL. Koch on yeast cells. *Soil Biol Biochem*, 30(8/9), 1998 1137-1145.
21. Moody SA, Briones MJI, Pierce TG and Dighton J. Selective consumption of decomposing wheat straw by earthworms. *Soil Biol Biochem*, 28, 1995, 533-537.
22. Edwards CA. Earthworm ecology (2^d edition), CRC Press LLC, Boca Raton, FL, USA. 2004, 12-23.
23. Horn M A, Schramma A and Draka H. The earthworm gut, An ideal habitat for ingested N₂O producing microorganisms. *Applied & Environmental Microbiology*, 69, 2003, 1662-1669.
24. Edwards CA and Bohlen PJ. Biology and Ecology of earthworms (3d edn), Chapman and Hall, London, 1996, 35-55.
25. Suthar S Microbial decomposition efficiencies of monoculture and polyculture vermireactors based on epigic and anecic earthworms. *World J Microbial Technology*, 24, 2008, 1471-1479.
26. Giraddi RS, Meenatchi R, Suresh B, Biradar MD and Biradar DP Standardisation of method for genomic DNA extraction in earthworms. *Karnataka J Agricultural Sci*, 22(4), 2009, 918-920.
27. Thakuria D, Olaf Schmidt, Dillon Finan, Damian Egan and Fiona M Doohan. Gut wall bacteria of earthworms, a natural selection process. *ISMEJ*, 4, 2010, 357-366.
28. Brown GG. How do earthworms affect microfloral and faunal community diversity?. *Plant Soil*, 170, 1995, 209-231.
29. Tereshchenko NN and Naplekova NN. Influence of different ecological groups of earthworms on the intensity of nitrogen fixation. *Biol Bull Rus Acad Sci*, 29, 2002 628-632.
30. Barois I, Villemin G, Lavelle P and Toutain F. Transformation of the soil structure through *Pontoscolex corethrurus* (Oligochaeta) intestinal tract. *Geoderma*, 56, 1993, 57-66.
31. Shipitalo MJ and Protz R. Chemistry and micromorphology of aggregation in earthworms casts. *Geoderma*, 45, 1989 357-374.
32. Nechitaylo TY, Yakimov MM, Godinho M, Timmis KN, Belogolova E, Byzov BA, Kurakov AV, Jones DL and Golyshin PN. Effect of the Earthworms *Lumbricus terrestris* and *Aporrectodea caliginosa* on bacterial diversity in Soil. *Microb Ecol*, 59, 2010 574-587.
33. Byzov BA, Khomyakor NV, Kharin SA and Kurakar AV. Cate of soil bacteria and fungi in the gut of earthworm. *Eur J Soil*, 43, 2007, 146-156.
34. Singleton DR, Hendrix PF, Coleman DC and Whitman WB. Identification of uncultured bacteria tightly associated with the intestine of the earthworm *Lumbricus rubellus* (Lumbricidae, Oligochaeta). *Soil Biol Biochem*, 35, 2003, 1547-1555.
35. Kritufek V, Ravasz K and Pizl V. Changes in densities of bacterial and microfungi during gut transit in *Lumbricus rubellus* and *Aporrectodea caliginosa* (Oligochaeta, Lumbricidae). *Soil Biol Biochem*, 24, 1992, 1499-1500.
36. Karsten GR and Drake HL. Comparative assessment of the aerobic and anaerobic microfloras of earthworm guts and forest soils. *Appl Environ Microbiol*, 61, 1995, 1039-1044.
37. Jolly JM, Lappin-Swtt HM, Anderson JM, Clegg CD. Scanning Electron Microscopy of the gut microflora of two earthworms, *Lumbricus terrestris* and *Octolcsion cyaneum*. *Microbial Ecology*, 26, 1993, 235-245.
38. Karsten GR and Drake HL. Denitrifying bacteria in the earthworm gastrointestinal tract and in vivo emission of nitrous oxide (N₂O) earthworms. *Appl Environ Microbiol*, 63, 1997, 1878-1882.
39. Singleton DR, Hendrix PF, Coleman DC and Whitman WB. Erratum to, identification of uncultured bacteria tightly associated with the intestine of the earthworm *Lumbricus rubellus* (Lumbricidae, Oligochaeta). *Soil Biol Biochem*, 36, 2004, 873.
40. Santiago A. Estudio de flora bacteriana intestinal de *Onychochaeta borincana* (Oligochaeta, alossoscolecidae) en Puerto Rico, MS Thesis, University of Puerto Rico, Mayaguez, 1995.
41. Satchell JE. *Lumbricidae*. In, Burges A, Raw F(ed) Soil biology. Academic press, London. 1967, 259-322.
42. Hendriksen NB. Gut load and food retention time in the earthworms *Lumbricus festivus* and *Lumbricus castaneus*. *Biol Fertil Soils*, 11, 1991, 170-173.
43. Matthies C, Griebhammev A, Schmittroth M and Drake HL. Evidence for involvement of gut-associated denitrifying bacteria in emission of nitrous oxide (N₂O) by earthworms obtained from garden and forest soils. *Appl Environ Microbiol*, 65, 1999, 3599-3604.
44. Elliott P W, Knight D and Anderson J M Variables controlling denitrification from earthworm casts and soil in permanent pastures. *Biol. Fertil. Soils*. 11, 1991, 24-29.



45. Svensson BH, Bostrom U and Klemedtson L. Potential for higher rates of denitrification in earthworm casts than in the surrounding soil. *Biol Fertil Soils*, 2, 1986, 147-149.
46. Bonkowski M, Griffiths BS and Ritz K. Food preference of earthworms for soil fungi. *Pedobiologia*, 44, 2000, 666-676.
47. Mindermann GS and Daniels L. Colonization of newly leaves by microorganisms, in, progress in soil biology Braunschweig, 1967, 3-9.
48. Pokarzhevskii AD, Sikora G and Gordienko SA. Amino-acid resources in the food of saprophages. *Dokl Acad Nauk SSSR*. 277, 1984, 253-256.
49. Sampedro L, Jeannotte RS and Whalen JK. Trophic transfer of fatty acids from the gut microbiota to the earthworm *Lumbricus terrestris* L in abstracts the 8th Inter. Symposium on Earthworm Ecology, 4-9 September, Krakow, Poland. 2006, 134.
50. Kulinska D. The effect of earthworms on the soil microflora. *Acta Microbiologica Polonica*, 10, 1961, 339-346.
51. Vaclav Pi L and Alena Novakova. Interactions between microfungi and *Eisenia andrei* (Oligochaeta) during cattle manure vermicomposting. *Pedobiologia*. 47(5-6), 2003, 895-899.
52. Binet F, Fayolle L, Pussard M, Crawford JJ and Traina SJ. Significance of earthworms in stimulating soil microbial activity. *Biol Fertil Soils*, 27, 1998, 79-84.
53. Morgan M and Burrows I Earthworms/Microorganisms interaction. Rothamsted Experimental Station, USA, 1982.

