



## ANTERIOR ESTHETIC GINGIVAL DEPIGMENTATION WITH DIODE LASER-A CASE REPORT

**Kamal Sagar<sup>1\*</sup>, Ritika Arora<sup>2</sup>, Vijay Kumar<sup>3</sup>, Pooja Patel<sup>4</sup>**

Demostartor<sup>1</sup>, Reader<sup>2</sup>, Department of Periodontics, Post Graduate Institute of Dental Sciences, Rohtak, Haryana, India.

Corresponding Author:- **Dr. Kamal Sagar**  
E-mail: [dr.kamalsagar@gmail.com](mailto:dr.kamalsagar@gmail.com)

<p><b>Article Info</b></p> <p><i>Received 15/05/2015</i> <i>Revised 20/06/2015</i> <i>Accepted 25/06/2015</i></p> <p><b>Key words:</b> Gingival Hyperpigmentation, Diode Laser, Gingival depigmentation.</p>	<p><b>ABSTRACT</b></p> <p>Gingival hyperpigmentation is a major concern mainly for young female patients visiting the dentist. The patients with gummy smile and Hyperpigmentation in anterior gingival region are more concerned esthetically as compared to others. Most pigmentation is caused mainly by 5 pigments out of which melanin is the main culprit. Melanin hyper pigmentation usually does not present as a medical problem, but patients may complain about their unpleasant black gums. A case is reported here on the cosmetic correction of “hyperpigmented gums” using diode laser at 1.5W settings.</p>
--	---

### INTRODUCTION

Gingival hyperpigmentation is increased pigmentation beyond the normally expected degree of the oral mucosa [1]. Melanin is derived from the Greek word “melas,” which means black [2]. It is an endogenous pigment produced by the melanocytes present in the basal and suprabasal layers of the epithelium [3]. The most common cause of hyperpigmentation is physiologic or ethnic as a result of excessive melanin deposition by melanocytes, which in turn depends on the activity of enzyme tyrosinase [4,5]. The pigmentation is mainly caused by five primary pigments including Melanin, melanoid, reduced hemoglobin, bilirubin, and iron [6-8]. The color of healthy gingiva is variable, ranging from a pale pink to a deep bluish purple hue. Between these limits of normalcy are a large number of pigmentation mosaics which depend primarily upon the intensity of melanogenesis, depth of epithelial cornification, and arrangement of gingival vascularity. Moreover, color variation may not be uniform and may exist as unilateral, bilateral, mottled, macular, or blotched, and may involve gingival papillae alone or extend throughout the gingiva on to other soft tissues [9,10]. It is a common esthetic

concern, aggravated in patients with a gummy smile or excessive gingival display. A periodontal plastic surgical procedure called the gingival depigmentation is performed whereby the gingival hyperpigmentation is removed or reduced by various techniques. This case report presents a case of gingival hyperpigmentation which was successfully treated by using diode laser.

### Case Report

A 27 - year old female patient visited in the outpatient department of maulana azad institute of dental sciences with the chief complaint of “black” coloured gums (Fig-1). Her oral examination revealed that she had deeply pigmented gingiva from right first premolar to left first premolar in both upper and lower arches. The patient requested for any kind of esthetic treatment which could make her “black” coloured gums look better. Depigmentation of maxillary arch with diode laser was planned in the first sitting. The entire procedure was explained to the patient and written consent was obtained. A complete medical, family and personal history with other routine investigations were carried out to rule



out any contraindication for surgery or any other hormonal disorder related to hyperpigmentation. Local anesthesia was infiltrated in the maxillary anterior region from premolar to premolar (Lignocaine with adrenaline in the ratio 1:100000 by weight). The diode laser (890 nm) was set at 1.5 W power, continuous contact mode to remove hyperpigmented areas (Fig-2). The laser beam was guided in a “brush-stroke” pattern until the entire area was free of

pigmentation. The area was irrigated with 1% normal saline in between the procedure. After 2 weeks of upper arch surgery, lower arch was also treated in the same fashion (Fig-3). No post-Operative discomfort, swelling was noticed. Twelve weeks after the treatment, the surface was smooth and shiny, the pigmentation was absent and there were no visible traces of any re-pigmentation (Fig-4)

**Figure 1. Pre-Operative**



**Figure 2. Immediate Post-Op (Upper arch)**



**Figure 3. Immediate Post-Op (Lower arch)**



**Figure 4. Post-Op view after 12 weeks (Upper arch)**



## DISCUSSION

Gingival depigmentation is a treatment to remove the melanin hyperpigmentation. Several techniques have been used including mechanical, surgical, electrosurgical, cryosurgical, free gingival grafts, and lasers [11-18]. Scalpel surgery for depigmentation is a time-tested technique and remains the gold standard. Scalpel surgery causes unpleasant bleeding during and after the operation and it is necessary to cover the surgical site with periodontal dressing for 7 to 10 days [19].

However, laser depigmentation has gained increasing importance and interest in dentistry as it has many advantages over conventional surgical modalities [20]. Lasers were introduced in 1960 by Maiman and were brought into general practice by Dr William and Terry Myers [21]. Different lasers such as carbon dioxide (CO<sub>2</sub>) laser, Nd:YAG laser, semiconductor diode laser, argon laser, Er:YAG laser and Er,Cr:YSGG laser have been reported as effective, pleasant and reliable method with

minimal postoperative discomfort and faster wound healing for depigmentation procedure. The diode laser is a solid-state semiconductor laser that typically uses a combination of gallium (Ga), arsenide (Ar), and other elements, such as aluminum (Al) and indium (In), to change electrical energy into light energy.

Dental Laser energy has an affinity for different tissue components. The 980 nm diode laser has energy and wavelength characteristics that specially target the soft tissues [22]. It has an affinity for hemoglobin and melanin, therefore it is more efficient and better equipped to address deeper soft tissue problems. Diode lasers have high electrical to optical efficiency, are small lightweight and compact, hence portable and are quiet devices as compared to other solid state and gas lasers (such as Nd:YAG, KTP:YAG, Ho, YAG, argon, erbium family and CO<sub>2</sub>) [23].



## CONCLUSION

The diode laser is a minimally invasive treatment option for the elimination of unesthetic gingival hyperpigmentation. There are many advantages of using diode laser in periodontal plastic surgeries like minimal

bleeding, fast healing and more control over the procedure.

The results in this case were excellent at 3-months follow-up period. There was no evidence of repigmentation of the gingiva resulting in improved esthetics.

## REFERENCES

1. Rashmi Hegde, Padhye Ashvini, Sumanth S, Jain, A. Sanjay, Thukral Naresh. (2013). Comparison of Surgical Stripping, Erbium-Doped, Yttrium, Aluminum, and Garnet Laser, and Carbon Dioxide Laser Techniques for Gingival Depigmentation, A Clinical and Histologic Study. *J Periodontol*, 84, 738-748.
2. Dummett CO (1945) Clinical observation on pigment variations in healthy oral tissues in the Negro. *J Dent Res*, 24, 7-13.
3. Dummett CO, Sakamura J, Barrens G. (1981). Attitudes towards normal pigmentation of the oral mucosa. *Quintessence Int*, 10, 115-122.
4. Newman MG, Takei HH, Klokkevold PR, Carranza FA. (2006). The gingiva. In, Carranza's Clinical Periodontology 10<sup>th</sup> ed. W.B. Saunders, Philadelphia, 46-66.
5. Dummett CO, Barends G. (1967). Pigmentation of the oral tissues, A review of the literature. *J Periodontol*, 38, 369-378.
6. Cicek Y, Ertas U. (2003). Normal and pathological pigmentation of oral mucous membrane, a review. *J Contemp Dent Pract* 4, 76-86.
7. Dummett CO, Barends G. (1967). Pigmentation of the oral tissues, A review of the literature. *J Periodontol* 38, 369-78.
8. Carranza AC, Saglie FR. (1990). Clinical features of gingivitis In, Carranza FA. Glickman's clinical periodontology, WB Saunders Company, Philadelphia, 109-25.
9. Dummett CO, Barends G. (1971). Oromucosal pigmentation, An updated literary review. *J Periodontol*, 42, 726-36.
10. Dummett CO. (1979). Oral tissue color changes (I). *Quintessence Int* 10, 39-45
11. Hirschfeld I, Hirschfeld L. (1951). Oral pigmentation and a method of removing it. *Oral Surg Oral Med Oral Pathol*, 4, 1012-1016.
12. Gnanasekhar JD, al-Duwairi YS. (1998). Electrosurgery in dentistry. *Quintessence Int* 29, 649-654.
13. Tal H, Landsberg J, Kozlovsky A. (1987). Cryosurgical depigmentation of the gingiva. A case report. *J Clin Periodontol*, 14, 614-617.
14. Tamizi M, Taheri M. (1996). Treatment of severe physiologic gingival pigmentation with free gingival autograft. *Quintessence Int*, 27, 555-558.
15. Novaes AB Jr, Pontes CC, Souza SL, Grisi MF, Taba M Jr. (2002). The use of acellular dermal matrix allograft for the elimination of gingival melanin pigmentation, Case presentation with 2 years of follow-up. *Pract Proced Aesthet Dent*, 14, 619-623
16. Nakamura Y, Hossain M, Hirayama K, Matsumoto K. (1999). A clinical study on the removal of gingival melanin pigmentation with the CO(2) laser. *Lasers Surg Med*, 25, 140-147.
17. Atsawasuwan P, Greethong K, Nimmanon V. (2000). Treatment of gingival hyperpigmentation for esthetic purposes by Nd, YAG laser, Report of 4 cases. *J Periodontol*, 71, 315-321.
18. Azzeh MM. (2007). Treatment of gingival hyperpigmentation by erbium-doped, yttrium, aluminum, and garnet laser for esthetic purposes. *J Periodontol*, 78, 177-184.
19. Kanakamedala AK, Geetha A, et al. (2010). Management of gingival hyperpigmentation by surgical scalpel technique. *J Clin Diagnos Res*, 4, 2341-46
20. Hibst R, Keller U. (1989). Experimental studies of the application of the Er, YAG laser on dental hard substances, I. Measurement of the ablation rate. *Lasers Surg Med*, 9, 338-344.
21. Doshi Y, Khandge N, Byakod G, Patil P. (2012). Management of gingival pigmentation with diode laser, Is it a predictive tool? *Int J Laser Dent*, 2(1), 29-32.
22. Gupta G. (2011). Management of gingival hyperpigmentation by semiconductor diode laser. *J Cutan Aesthet Surg*, 4(3), 208-10.
23. Mathews MA. (2011). Diode lasers, A versatile clinical tool (a technical and clinical review). *Intl J Laser Dent*, 1(1), 9-15.

