



ACCESSORY BRACHIALIS MUSCLE - A CASE REPORT

Ansari Kainat Khalid Akhtar

Department of Anatomy, K.J. Somaiya Medical College, Somaiya Ayurvihar, Eastern Express Highway, Sion, Mumbai-400 022, Maharashtra, India.

Corresponding Author:- **Ansari Kainat Khalid Akhtar**

E-mail: kansari4@yahoo.com

<p>Article Info <i>Received 15/01/2016</i> <i>Revised 27/02/2016</i> <i>Accepted 12/03/2016</i></p> <p>Key words: Accessory Brachialis Muscle, Median Nerve, Brachial Artery, Variation, Nerve Compression.</p>	<p>ABSTRACT</p> <p>During routine dissection for first M.B.B.S. students on 65 years donated embalmed male cadaver in the Department of Anatomy, K. J. Somaiya Medical College, we encountered an accessory brachialis muscle taking origin from the anteromedial surface of shaft and medial supracondylar ridge of lower end of humerus and inserted into the medial epicondyle of humerus. The median nerve and brachial artery pierced the muscle. The role of additional muscles in compression syndrome is a well-known phenomenon. Medical fraternity including orthopaedicians and neurologists need to be aware of such variations when dealing with upper limb injuries or operations around the elbow joint.</p>
---	--

INTRODUCTION

Brachialis arises from the lower half of the front of the humerus and from the medial intermuscular septa, converge to a thick, broad tendon which is attached to the ulnar tuberosity and to a rough impression on the anterior aspect of the coronoid process. Brachialis may be divided into two or more parts. It may be fused with brachioradialis, pronator teres or biceps. In some cases it sends a tendinous slip to the radius or bicipital aponeurosis [1]. Surgeons and orthopedists dealing with cases of road traffic accidents need to be aware of such variations as dealing with such cases may not only be difficult, but a nightmare too.

CASE REPORT

During routine dissection for first M.B.B.S. students on 65 years donated embalmed male cadaver in the Department of Anatomy, K. J. Somaiya Medical College, we encountered an accessory brachialis muscle taking origin from the anteromedial surface of shaft and medial supracondylar ridge of lower end of humerus and inserted into the medial epicondyle of humerus. The median nerve and brachial artery pierced the muscle.



DISCUSSION

Several cases on the presence of accessory brachialis muscle in the arm have been reported. Dharap observed an unusual muscle that passed obliquely from the middle of the humerus anterior to the median nerve and brachial artery, forming a tunnel for them, before inserting with the common origin of the forearm flexor muscles [2]. An accessory brachialis (AcBr) muscle originating from mid shaft of humerus and medial intermuscular septum has been reported. During its course medially, toward the elbow, the AcBr crossed both the brachial artery and the median nerve. The distal tendon split to surround the median nerve before inserting into the brachialis tendon and the common tendon of the antebrachial flexor compartment muscles [3]. A variant muscle arising from the medial border of brachialis muscle has been described which after bridging the median nerve, the brachial artery and vein; was fusing with the medial intermuscular septum. The muscle was innervated by musculocutaneous nerve [4]. Few fleshy fibers of brachialis arising from the distal third of the muscle and merging with superficial



flexors of the forearm and to the medial aspect of olecranon process of ulna have also been described [5]. The occurrence of an accessory brachialis muscle forming a fibro-muscular tunnel after blending with the medial intermuscular septum has been reported [6]. Although there are no symptoms most of the time, such structures have the potential to compress the median nerve with consequent functional impairment. Such accessory muscle slips may also compress the underlying arteries viz., brachial artery in the present case. Embryologically, the intrinsic muscles of the upper limb differentiate in situ, opposite the lower six cervical and upper two thoracic

segments, from the limb bud mesenchyme of the lateral plate mesoderm. The formation of muscular elements in the limbs takes place shortly after the skeletal elements begin to take shape. At a certain stage of development, the muscle primordia within the different layers of the arm fuse to form a single muscle mass [7]. Langman stated, however, that some muscle primordia disappear through cell death despite the fact that cells within them have differentiated to the point of containing myofilaments. Failure of muscle primordia to disappear during embryologic development may account for the presence of the accessory muscular bands reported in this case [8].



Clinical Significance

Compression of the median nerve and brachial artery by various types of structures leading to clinical neuro-vasculopathy has been reported [9]. On contraction, these muscles can compress the median and ulnar nerves, leading to further irritation of the nerves. Contraction of these muscles can compress both the brachial artery and brachial veins. Also, these muscles should not be mistaken for tumours on MR imaging of the arm [10].

CONCLUSION

The existence of such variation of the brachialis muscle should be kept in mind by the surgeons operating on patients with high median nerve palsy and brachial artery compression, by the orthopaedicians dealing with fracture of the humerus, the radiologists while doing radio-diagnostic procedures e.g. CT scan, MRI of the arm and angiographic studies and also by the physiotherapists. These accessory fibres of brachialis may be used as a transposition flap in deformities of infraclavicular and axillary areas and in post-mastectomy reconstruction. The accessory fibres of brachialis may prove significant and lead to confusion during surgical procedures or cause compression of neurovascular structures.

REFERENCES

1. Standing S. (2005). Gray's Anatomy, 39th Ed, London, Elsevier Churchill Livingstone, 855.
2. Dharap AS. (1994). An anomalous muscle in the distal half of the arm. *Surg Radiol Anat*, 16, 97-99.
3. Loukas M, Louis RG Jr, South G, Alsheik E, Christopherson C. (2006). A case of an accessory brachialis muscle. *Clin Anat*, 19, 550-553.

ACKNOWLEDGEMENT

The author wishes to convey sincere thanks to Dean Dr. Geeta Niyogi Madam for her valuable help, support and inspiration. The author is also thankful to the Head of Department Dr. Sawant and all staff members of Department of Anatomy. The author also acknowledges the immense help received from the scholars whose articles are cited and included in references of this manuscript. The authors are also grateful to authors / editors / publishers of all those articles, journals and books from where the literature for this article has been reviewed and discussed.

CONFLICT OF INTEREST

The authors declare that they have no conflict of interest.

STATEMENT OF HUMAN AND ANIMAL RIGHTS

All procedures performed in human participants were in accordance with the ethical standards of the institutional research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards. This article does not contain any studies with animals performed by any of the authors.



4. Paraskevas G, Natsis K, Loannidis O, Papaziogas B, Kitsoulis P, Spanidou S. (2008). Accessory muscles in the lower part of the anterior compartment of the arm that may entrap neurovascular elements. *Clin Anat*, 21, 246–251.
5. George BM, Nayak SB. (2008). Median nerve and brachial artery entrapment in the abnormal brachialis muscle – a case report. *Neuroanatomy*, 7, 41–42.
6. Vadgaonkar R, Rai R, Ranade AV, Nayak SR, Pai MM, Lakshmi R. (2008). A case report on accessory brachialis muscle. *Rom J Morphol Embryol*, 49, 581–583.
7. Arey LB. *Developmental Anatomy*. (1960). In, *A Textbook and Laboratory Manual of Embryology*. 6th Ed., Philadelphia, WB Saunders Company, 434–435.
8. Langman J. *Medical Embryology*. 2nd Ed, (1969), Baltimore, Williams and Wilkins, 143–146.
9. Gessini L, Jandolo B, Pietrangeli A. (1983). Entrapment neuropathies of the median nerve at and above the elbow. *Surg Neurol*, 19, 112–116.
10. Sharadkumar Sawant. (2015). Study of variant distal attachment of brachialis muscle. *International Journal of Pharmacology Research*, 5(1), 44-47.

