



BODY MASS INDEX ON LUNG FUNCTION IN HEALTHY MALE AND FEMALE ADULTS

Arifath Gopinath^{1*}, Lalitha Shanmugam²

¹Associate Professor of of Physiology, Melmaruvathur Adhiparasakthi Institute of Medical Sciences and Research Institute, Melmaruvathur, Tamilnadu, India

² Assistant Professor of of Physiology, Melmaruvathur Adhiparasakthi Institute of Medical Sciences and Research Institute, Melmaruvathur, Tamilnadu, India.

ABSTRACT

Obesity is becoming one of the serious public health problems of modern world with rapidly changing lifestyles involving consumption of high calorie foods with decreased physical activities. Spirometry is the initial screening tool for pulmonary diseases. The aim of this study was to evaluate the prevalence of deranged BMI, pulmonary function tests and correlation between BMI and pulmonary function test. Our aim is to find association and correlation between BMI and pulmonary functions parameters. Increasing BMI is associated with decrease in lung functions in healthy population of either gender. Regular pulmonary function monitoring may assess adverse effects of obesity on pulmonary functions. Hence, proper and timely advice regarding lifestyle modification to obese subjects will prevent unwanted complications of obesity.

Keywords:- Body Mass Index (BMI), Pulmonary Function Test (PFT), Forced Vital Capacity (FVC), Spirometry.

Access this article online

Journal Home Page

www.mcmed.us/journal/abs

Quick Response code



Received:29.09.2018

Revised:17.10.2018

Accepted:19.11.2018

INTRODUCTION

Lung function is a significant predictor of future morbidity and mortality in the general population.[1] Maintaining good lung function across adult life is important to prevent chronic respiratory diseases, which nowadays represent a serious public health problem around the world.[2] There is consistent evidence showing that overweight, obesity and weight gain in adulthood are detrimental to lung function, as described by the forced vital capacity (FVC) and/or forced expiratory volume in 1 s (FEV1). Previous population-based and occupational cohort studies have shown that excessive weight gain in adulthood is associated with lower lung function levels and with an increased rate of lung function decline independently of age and smoking status.[3-8]. This precludes a more comprehensive understanding of the role of weight change on lung function during adulthood and older life and supports the

need for further studies with longer follow-up periods extending into late adulthood. Understanding the effects of weight changes on lung function during adult life is of utmost importance given the epidemic levels of overweight and obesity globally.[9] BMI has also shown its importance in the prediction of various disease conditions, the most important being the predictor of cardiovascular risks.[10] However, the link between BMI and lung function test is yet to be explored in detail.[11] Previous studies have well established the association of age, sex, and anthropometric indices with pulmonary functions.[12-16]

Spirometry is now widely acceptable test to assess pulmonary functions. They are considered as the initial screening tool for pulmonary diseases and are easy to conduct by using equipment that is available in all pulmonary functions laboratories.

Therefore, our study aimed to assess the effects of BMI on spirometric test parameters among young healthy adults. Our aim is to find association and correlation between BMI and pulmonary functions parameters.

Materials and methods:

This is a cross sectional study conducted in Melmaruvathur Adhiparasakthi Institute of Medical Sciences and Research Institute, Melmaruvathur. After approval from institutional ethics committee, total 70 adult subjects in the age group of 18 to 75 years were selected. Both males and females were included. The subjects selected were healthy volunteers, hospital staff and relatives of patients. Informed written consent was taken from all the participants. Each individual underwent a thorough medical evaluation including medical history and complete physical examinations. Current smokers, ex-smokers and tobacco users in any form like chewing, snuffing or water pipe were excluded from this study. Subjects with pre-existing pulmonary diseases like tuberculosis, bronchial asthma, COPD etc or having systemic conditions like diabetes mellitus, hypertension, as cites etc were also excluded. Patients having history of recent cardiac or thoracic surgery or having chest deformities or serious medical conditions were also not considered.

Measurement of anthropometric parameters:

Standing height of subject was recorded to the nearest one centimet reusing Bio-plus stadiometer, without footwear, with heels together and heels, calf, buttocks and preferably back touching the stadiometer. The weight was measured by eagle's weighing machine to the nearest 0.1 kg, in standing position; subjects were wearing light clothes and were bare footed. Body mass index (BMI) was calculated by using Quetelet's index. ($BMI = \text{Weight in kg} / \text{Height in m}^2$). [17] Depending on the BMI values, the subjects were divided into obese, overweight and non obese groups as per WHO

classification system. The first group consisted of non-obese (normal body weight) subjects with BMI of 18 to 24.9 kg/m², second group consisted of BMI of 25 to 29.9 kg/m² and obese subjects with BMI of 30 kg/m² and above.

Measurement of Pulmonary Function Tests:

The pulmonary function parameters were measured by using computerized body plethysmograph (Medgraphics – Platinum Elite Sepies™ Plethysmograph) machine in the Department of Pulmonary Medicine. The tests were conducted according to the American

Thoracic Society/ European Respiratory Society (ATS/ERS) task force guidelines. [18] The instrument was calibrated daily. All the subjects were instructed to avoid tea, coffee and other stimulants and to report after a light breakfast. The test was explained and demonstrated to the subjects. After a rest for 5–10 minutes, the test was carried out in the sitting position, wearing a nose clip. The best of the three acceptable results were selected. Parameters recorded were; Forced vital capacity (FVC) in litres, Force expiratory volume in one second (FEV₁) in litres, FEV₁/FVC, Peak expiratory flow rate (PEFR) in litre per second, Forced expiratory flow rate 25-75%. All spirometric parameters were considered as a percentage of predicted on reported height and age. [19]

Statistical Analysis:

The detailed data was entered into the Microsoft excel sheet and subsequently analyzed statistically spss-11.5. Obtained Values were reported as Mean \pm S.D. Comparison of lung functions in males and females was done using student's 't' test. Comparison of lung functions in normal, overweight and obese subjects was done using ANOVA test. Pearson's correlation coefficient was applied to determine the correlation between pulmonary functions parameters and BMI. Significance level was set at $p < 0.05$ and considered as significant.

Results:

Table 1: Baseline characteristics of study population according to gender

Variables	Male	Female	p-value
Age(Years)	47.12 \pm 15.16	45.44 \pm 15.68	0.51
Height (cm)	153.08 \pm 06.08	164.45 \pm 09.84	<0.0001*
Weight(Kg)	64.34 \pm 16.78	71.55 \pm 17.78	0.015*
BMI (Kg/m ²)	27.67 \pm 06.98	26.84 \pm 09.08	0.52
FVC(%)	88.50 \pm 11.72	88.81 \pm 12.89	0.86
FEV ₁ (%)	76.59 \pm 12.30	76.30 \pm 11.63	0.87
FEV ₁ / FVC	86.15 \pm 10.66	85.87 \pm 10.87	0.21
EFF25-75%	87.43 \pm 7.41	86.58 \pm 4.64	0.82
PEFR(%)	249.58 \pm 46.09	352.9 \pm 53.75	<0.0001*

Table 2: Comparison of characteristics of various BMI categories in males

Variables	BMI category (kg/m ²)			p value
	Normal (n=34)	Overweight (n=26)	Obese (n=10)	
Age (Years)	49.14 ± 16.65	46.42 ± 14.47	42.60 ± 11.63	0.42
Height (cm)	164.09 ± 7.25	165.06 ± 4.66	159.23 ± 20.36	0.11
Weight (Kg)	59.55 ± 6.65	74.54 ± 5.40	101.69 ± 21.07	<0.0001*
BMI (Kg/m ²)	21.79 ± 1.49	27.01 ± 1.39	41.94 ± 15.17	<0.0001*
FVC (%)	80.87 ± 12.45	79.23 ± 11.37	82.21 ± 12.38	0.10
FEV1 (%)	77.70 ± 11.32	72.50 ± 9.25	65.42 ± 12.53	0.06*
FEV1 / FVC	96.21 ± 11.08	88.48 ± 8.37	83.42 ± 12.38	0.04*
EFF 25-75%	91.19 ± 26.08	84.34 ± 23.73	76.42 ± 21.35	0.21
PEFR (%)	360.59 ± 102.55	302.67 ± 86.18	252.58 ± 105.37	0.020*

Table 3: Comparison of characteristics of various BMI categories in females

Variables	BMI category (kg/m ²)			p value
	Normal (n=28)	Overweight (n=18)	Obese (n=24)	
Age (Years)	44.53 ± 17.09	42.09 ± 13.82	48.46 ± 15.79	0.42
Height (cm)	153.88 ± 06.20	152.83 ± 06.44	152.29 ± 5.83	0.62
Weight (Kg)	51.14 ± 07.09	63.90 ± 06.93	80.04 ± 16.18	<0.0001*
BMI (Kg/m ²)	21.46 ± 01.75	27.24 ± 01.00	35.20 ± 05.67	<0.0001*
FVC (%)	87.57 ± 12.18	88.09 ± 12.88	89.21 ± 11.38	0.91
FEV1 (%)	76.25 ± 12.75	75.47 ± 11.88	73.42 ± 12.53	0.42
FEV1 / FVC	87.32 ± 10.32	85.21 ± 08.37	82.83 ± 12.38	0.04
EFF 25-75%	91.10 ± 14.49	94.34 ± 23.73	78.54 ± 31.40	0.013*
PEFR (%)	248.48 ± 67.74	208.67 ± 86.18	161.49 ± 75.51	0.015*

Table 4: Correlation between BMI and lung function parameters in different BMI categories in both genders

BMI Category	Parameter	Male		Female	
		r ²	p value	r ²	p value
Normal	FVC (%)	0.01	0.46	0.0096	0.91
	FEV1 (%)	0.044	0.13	0.007	0.78
	FEV1 / FVC	0.008	0.49	0.036	0.26
	EFF 25%-75%	0.001	0.69	0.007	0.77
	PEFR (%)	0.013	0.39	0.009	0.92
Overweight	FVC(%)	0.119	0.02	0.046	0.37
	FEV1(%)	0.929	< 0.0001*	0.17	0.10
	FEV1/ FVC	0.003	0.65	0.19	0.09
	EFF25%-75%	0.039	0.22	0.21	0.07
	PEFR(%)	0.003	0.67	0.03	0.42
Obese	FVC(%)	0.249	0.0145	0.5812	< 0.0001*
	FEV1(%)	0.179	0.0881	0.5953	< 0.0001*
	FEV1/ FVC	0.419	0.0302*	0.9631	< 0.0001*
	EFF25%-75%	0.009	0.6277	0.03543	0.30
	PEFR(%)	0.059	0.377	0.02624	0.35

Discussion:

Obesity is one of the major health hazards across the world. It can lead to various clinical complications such as diabetes, vascular diseases, osteoarthritis, etc. But less emphasis has been given on the effect of obesity on respiratory system.[20] In this study an attempt was made to find out whether there is an increased risk of

respiratory problems in overweight and obese individuals. Pulmonary function tests are generally related to body size and age, where height is a proxy for chest size, and age reflects maturity.[21] Because of this reason every individual has different range of normal values.

In present study it was observed that in overweight and obese male subjects FEV1, FEV /FVC

and PEFR values were significantly lower. This can be explained on the basis of fact that with increase in obesity, respiratory and airway resistance increases.[22] Some investigators also observed similar findings in their studies.[23,24] But Costa et al reported nonsignificant difference of lung functions among obese and non-obese females.[25]. In overweight and obese female subjects FEF_{25-75%} and PEFR values were observed to be significantly lower. This shows obstructive pattern of smaller airways. Due to increased fat deposition on the chest wall, expansion of thoracic cavity is affected.[26] Also fat deposition in abdominal cavity shifts diaphragm upwards decreasing lung functions further and increasing work of breathing.

We observed significant negative correlation between BMI and FEV₁ in overweight males. Obese females showed significant negative correlation between BMI and FEV₁, FVC, FEV₁/FVC; while males showed significant negative correlation between BMI and FEV₁/FVC₁. Similar results were also observed by many other studies.[27-29] It may be because of fat accumulation around ribs, abdomen and diaphragm which causes restricted movements of ribs, reducing lung volume and decreasing respiratory compliance.[30].

All the parameters of pulmonary function tests were negatively correlated with Body mass index (BMI) in rural population. The present findings were not supported by the study done by Piyali et al. [31] The possible cause of the difference between two studies may be age factor and mild COPD for both sexes in their study. Therefore, it can be said that obesity has significant

impact on respiratory problems. Excess of abdominal fat may restrict the diaphragmatic movement which leads to a decrease in pulmonary function. This study suggested significant impairment of pulmonary functions in overweight and obese population due to limited expansion of thoracic cavity which leads to possibility of small airway diseases. The lung functions might be improved by weight loss. According to WHO, BMI is the gold standard to classify obesity at present time. But BMI does not take an account of body fat distribution. Thoracic and abdominal fat have direct effects on movement of diaphragm rather than fat in hips and thighs. Hence, along with BMI, other indicators of fat distribution should also be considered.

Conclusion:

Moderate and high weight gain over 20 years was associated with accelerated lung function decline, while weight loss was related to its attenuation. Control of weight gain is important for maintaining good lung function in adult life. No significant difference was detected in spirometry parameters between obese/overweight and normal weight school children in our study. Increasing BMI is associated with decrease in lung functions in healthy population of either gender. Regular pulmonary function monitoring may assess adverse effects of obesity on pulmonary functions. Hence, proper and timely advice regarding lifestyle modification to obese subjects will prevent unwanted complications of obesity.

REFERENCES

1. Young RP, Hopkins R, Eaton TE. Forced expiratory volume in one second: not just a lung function test but a marker of premature death from all causes. *Eur Respir J* 2007;30:616–22.
2. Young RP, J. Hopkins R. Primary and secondary prevention of chronic obstructive pulmonary disease: where to next? *Am J Respir Crit Care Med* 2014;190:839–40.
3. Wang M-L, McCabe L, Peterson EL, et al. Weight gain and longitudinal changes in lung function in steel workers. *Chest* 1997;111:1526–32.
4. Bottai M, Pistelli F, Di Pede F, et al. Longitudinal changes of body mass index, spirometry and diffusion in a general population. *Eur Respir J* 2002;20:665–73.
5. Carey IM, Cook DG, Strachan DP. The effects of adiposity and weight change on forced expiratory volume decline in a longitudinal study of adults. *Int J Obes* 1999;23:979–85.
6. Chen Y, Horne SL, Dosman JA. Body weight and weight gain related to pulmonary function decline in adults: a six year follow up study. *Thorax* 1993;48:375–80.
7. Chinn DJ, Cotes JE, Reed JW. Longitudinal effects of change in body mass on measurements of ventilatory capacity. *Thorax* 1996;51:699–704.
8. Chinn S, Jarvis D, Melotti R, et al. Smoking cessation, lung function, and weight gain: a follow-up study. *Lancet* 2005;365:1629–35.
9. The World Health Organization. Obesity and overweight. Available: <http://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight> [Accessed 22 Aug 2018].
10. Mishra S, Banerjee S, Sengupta TK, Behera AA, Manjareeka M, Mishra J. Association of diet and anthropometric measures as cardiovascular modifiable risk factors in young adults. *J Basic Clin Physiol Pharmacol* 2013;1–8.]
11. Jones RL, Nzekwu MM. The effects of body mass index on lung volumes. *Chest* 2006;130:827–33.

12. Mishra J, Mishra S, Satpathy S, Manjareeka M, Nayak PK, Mohanty P. Variations in PEFR among males and females with respect to anthropometric parameters. *IOSR J Dent Med Sci* 2013;5:47-50.
13. Choudhuri D, Choudhuri S. Effect of gender and body mass index on pulmonary function tests in adolescents of tribal population of a North Eastern State of India. *Indian J Physiol Pharmacol* 2014;58:170-3.
14. Ajmani S, Anupama N, Nayanatara AK, Sharma VM, Ganaraja B, Pai SR. Effect of abdominal fat on dynamic lung function tests. *Int J Biomed Adv Res* 2012;3:632-6.
15. Manjareeka M, Mishra J, Padhi RK, Mishra S, Nanda S. Peak expiratory flow rate as a function of anthropometric variables in tribal school children. *Int J Physiol* 2014;2:4-8.
16. Manjareeka M, Mishra J, Nanda S, Mishra S, Mishra JP. Assessment of peak expiratory flow rate in preadolescent children of sub-tribal communities in Odisha. *Int J Clin Exp Physiol* 2014;2:120-4.
17. Park K. Obesity. In: Park's textbook of preventive and social medicine. 21st ed. Jabalpur, India: Banarsidas Bhanot; 2011. p. 366-70.
18. Miller MR, Hankinson J, Brusasco V. Standardization of spirometry. *Eur Respir J* 2005;25:319-38.
19. Knudson RJ, Lebowitz M, Holberg CJ, Burrows B. Changes in the normal maximal expiratory flow-volume curve with aging. *Am Rev Resp Dis*. 1983;127:725-34.
20. Salome CM, King GG, Bernard N. "Physiology of obesity and effects on lung function." *J Appl Physiol*. 2010;108:206-11.
21. Stanojevic S, Wade A, Stocks J. "Reference values for lung function: past, present and future." *European Resp J*. 2010;36:12-19.
22. Carey IM, Cook DG, Strachan DP: The effects of adiposity and weight change on forced expiratory volume decline in a longitudinal study of adults. *Int J Obes Relat Metab Disord* 1999, 23(9):979-985.
23. Jones RL, Nzekwu MMU. The effect of body mass index on lung volumes. *Chest* 2006;130:827-33.
24. Al-Bader WR, Ramadan J, Nasr-Eldin A, Barac-Nieto M. Pulmonary ventilator functions and obesity in Kuwait. *Med Princ Pract*. 2008; 17:20-6.
25. Costa D, Barbalho MC, Miguel GPS, Forti EMP, Azevedo JLMC. The impact of obesity on pulmonary function in adult women. *Clinics* 2008; 63:719-24.
26. Poulain M, Doucet M, Major GC, Series F, Boulet LP, Tremblay A et al. The effect of obesity on chronic respiratory diseases: pathophysiology and therapeutic strategies. *CMAJ* 2006, 174:1293-1299.
27. Sin DD, Jones RL, Man SFP. Obesity is a risk factor for dyspnea but not for airflow obstruction. *Arch Intern Med*. 2002;162:1477-81.
28. Andrew J, Debbie B, Ali B. The Association of Body Mass Index with Airway Obstruction in Non-Asthmatics: Implications for the Inaccurate Differential Diagnosis of Asthma in Obesity. *Canadian Journal of Respiratory Therapy* 2011;47:2.
29. Rubinstein I, Zamel N, DuBarry L, Hoffstein V. Airflow limitation in morbidly obese, nonsmoking men. *Ann Intern Med*. 1990;112:828-32.
30. Salome CM, King GG, Bernard N. "Physiology of obesity and effects on lung function." *J Appl Physiol*. 2010;108:206-11.
31. Piyali S, Sumita M, Hiranmoy M, Dhara PC. Rural- Urban variation of pulmonary function in relation to age, sex and anthropometric variables among the Bengalee population. *Ergonomics for Rural Development*. 52-62.



Attribution-NonCommercial-NoDerivatives 4.0 International