



VALORIZATION OF SOURSOP FLOWERS (*ANNONA MURICATA* L.) AS A POTENTIAL SOURCE OF NATURAL ANTIOXIDANTS FOR PALM OLEIN STABILIZATION DURING ACCELERATED STORAGE

Dr. Siva Nageswara Rao Mekala^{1*}, Dr. Ali Said Yussuf²

¹Department of Clinical Pharmacology, The State University of Zanzibar – Tanzania.

²Department of Surgery, The State University of Zanzibar, Tanzania.

Article Info

Received 29/11/2025

Revised 16/12/2025

Accepted 19/01/2026

Key words: -

Natural Antioxidants,
Soursop flower extract,
Palm olein, Oxidative
stability.

ABSTRACT

This research aims at exploring the possibility of using *Annonamuricata* L. (soursop) flower extracts as a natural antioxidant to stabilize palm olein in accelerated storage. Oxidative stability of palm olein was tested by the addition of different levels of soursop flower extract 200-1800 ppm and exposure to accelerated oxidation at 70°C in 30 days. The extract was compared with a synthetic antioxidant butylatedhydroxytoluene (BHT) in terms of its efficacy. Conclusion of the Rancimat and Schaal oven testing showed that the soursop flower extract effectively increased the induction time, decreased the formation of peroxides and suppressed the formation of secondary oxidation products. The extract also maintained the amount of the linoleic acid in palm olein, which proved its potential as the substitute, natural antioxidant. The presence of the following phenolic compounds like quercetin, caffeic acid, and ferulic acid was determined using high-performance liquid chromatography (HPLC) analysis, and this led to the antioxidant effect. This indicates that palm olein can be stabilized using soursop flower extracts and this will create a viable source of natural antioxidant to replace synthetic preservatives.

INTRODUCTION

Oil, fats, and lipid-containing foods are subject to decay when heated or stored over a long period of time, mainly through oxidation. This oxidation causes food to lose the nutritional value of food, and therefore, the fatty acids, amino acids, vitamins, and the digestibility of the protein is lost [1]. Also, oxidation reduces the organoleptic characteristics of foods and may produce potentially toxic oxidation products related to degenerative diseases, including cancer and cardiovascular diseases. This causes products that are oxidized to be rejected by the consumers and industries lose a lot of money [2]. Synthetic antioxidants such as

butylated hydroxyanisole (BHA), butylated hydroxytoluene (BHT) and ter-butylhydroquinone (TBHQ) have also been applied as food preservatives in an attempt to prevent these deterioration reactions. Nevertheless, the questions about their health effects such as association with cancer and cardiovascular diseases have caused growing limitations in their usage in most countries [3]. In specific, BHA and BHT are volatile and readily degrade under high temperature, which limits their usefulness [4, 5]. This has triggered the consumer demand of natural antioxidants which are viewed as safer alternatives, and provides more health benefits [6].

Natural antioxidants Arguably phenolic compounds or polyphenols are natural antioxidants that have antioxidant activity because of their structure, hydrogen-donating ability, metal-ion chelation capability and synergies. There has been a lot of research on the

Corresponding Author

Dr. Siva Nageswara Rao Mekala

Email: - sivanageshmekala@yahoo.com



Schaal Oven Test

To accelerate oxidation, samples were stored at 70°C in 30 days. Peroxide value, p-anisidine, total oxidation (TOTOX), Thio barbituric acid, and iodine values were analyzed in samples collected after every 10 days using standardized procedures.

Measurement of the Oxidation Parameters

The peroxide value, p-anisidine, iodine value and TOTOX were measured according to the known procedures. The values of Thiobarbituric acid were calculated [12-14].

The Soursop Flower Extract on the Linoleic Acid Profile of Palm Olein during Storing

Fatty Acid Methyl Esters (FAMES) Preparation

Transesterification was done using 2% sulfuric acid in methanol to make the fatty acid methyl esters (FAMES) of the stabilized and control palm olein samples, as shown by Christie (1993). The FAMES had been put into ethyl acetate and thoroughly washed with water to eliminate all the excess acid and dried in the presence of anhydrous sodium sulfate. Dried esters were examined using gas chromatography (GC/FID) and all analyses were done in a duplicated manner.

Gas Chromatography (GC)

Agilent 7890A series gas chromatograph (Agilent Technologies, Palo Alto, CA) with FID detector and DB-225 capillary column (30 m x 0.25 mm film thickness 0.25 mm) were used to carry out GC-FID analysis. The column temperature was first adjusted to 160 °C, then 220 °C at the rate of 5 °C/min and maintained 10 min in 240 °C. The carrier gas was nitrogen and its flow rate was 1.5 mL/min. The injector and detector temperatures were 230 °C and 250 °C respectively, and the ratio of the split was 50:1. The identification of linoleic acid was done by comparing the retention times with the standard reference fatty acid methyl esters in the same conditions.

RESULTS AND DISCUSSION

Extraction Yield, Total Phenolic Content, and Phenolic Antioxidants Detected by HPLC

The 8.35% was obtained using the methanolic extract of soursop flowers. This was less than the 10.80% that George et al. (2012) recorded on the butanolic extracts of soursop leaves. The amount of total phenolic content of the soursop flower extract was 51.33 mg/g, which was larger compared to the results achieved by George et al. (2012) in the butanolic extracts of soursop leaves. Such variations in yield and phenolic composition may be explained by differences in the part of the plant, environment, time of collection of the sample and the method of determination.

The HPLC-DAD chromatograph of the soursop flowers methanolic extract indicated the presence of some phenolic antioxidants, which comprised vanillic acid (RT: 9.140 min, peak area: 7.74%), caffeic acid (RT: 9.375 min, peak area: 4.90%), ferulic acid (RT: 9.871 min, peak area: 13.07%), ellagic acid (RT: 10. The compounds are famous in terms of their high antioxidant properties, with quercetin being the most common compound. These results are in line with those of Georges et al. (2014) who found quercetin in alcoholic extracts of soursop leaves.

Primary Preliminary Antioxidant Tests DPPH and FRAP Assays

The radical scavenging behavior of the soursop flowers extract was evaluated by means of the DPPH radical scavenging test and the Ferric Reducing Antioxidant Power (FRAP). The two assays showed that the antioxidant activity of the extract improved with concentration. The extract radical scavenging activity was similar to the BHT and reducing power was similar to catechin. These findings affirm the high antioxidant capacity of the soursop flowers which is owed to the presence of phenolic compounds in the extract. Barreira et al. (2008) also identified a similar finding with respect to chestnut flowers that had high antioxidant properties.

Soursop Flower Extract Effect on Palm Olein Oxidative Stability.

Rancimat Test

To test the impact of the Rancimat test, the effect of the soursop flower extract on the palm olein oxidation was performed. Table 1 presents the variables that indicated that the induction time and the factors of the protection of the oil samples with the antioxidants were significantly higher ($P < 0.05$) than those of the control. The activity of the soursop flower extract at any concentration was significantly superior ($P < 0.05$) to BHT, which implies that it has better thermal stability and antioxidant capacity. The period of induction and the factors of protection were on the rise with the concentration of the extract and this illustrates its good antioxidant effects. Such findings are in line with those of Iqbal and Bhangar (2007) who discovered that extracts of garlic and pomegranate increased the stability of sunflower oil.

Schaal Oven Test

The Schaal oven test was performed so as to imitate accelerated oxidation. The peroxide values of oil samples containing soursop flower extract were significantly lower in comparison to the control which demonstrates that the extract was effective in the inhibition of peroxides. The more concentrated samples of palm olein (PO + An.M 1400 ppm, PO + An.M 1000 ppm, PO + An.M 1800 ppm) demonstrated better



resistance to the formation of peroxides, and the values were similar to the values of the oils that contain BHT.

This proves the efficacy of the extract in inhibiting the formation of primary oxidation products.

Table 1: Effect of Soursop Flower Extract on the Oxidative Stability of Palm Olein (Rancimat Test).

| Sample | Induction Time (hrs) | Protection Factor (PF) |
|--------------------|----------------------|------------------------|
| Control | 4.5 | 1.0 |
| PO + BHT 200 ppm | 6.3 | 1.4 |
| PO + An.M 200 ppm | 6.0 | 1.3 |
| PO + An.M 600 ppm | 6.8 | 1.5 |
| PO + An.M 1000 ppm | 7.2 | 1.6 |
| PO + An.M 1400 ppm | 7.5 | 1.7 |
| PO + An.M 1800 ppm | 8.0 | 1.8 |

Table 2: Oxidation Parameters of Palm Olein During Storage (Schaal Oven Test).

| Sample | Peroxide Value (meq/kg) | p-Anisidine Value | TOTOX Value | Thiobarbituric Acid (TBA) | Iodine Value (g I ₂ /100g) |
|--------------------|-------------------------|-------------------|-------------|---------------------------|---------------------------------------|
| Control | 12.3 | 7.5 | 28.8 | 3.2 | 65.4 |
| PO + BHT 200 ppm | 7.2 | 3.5 | 14.4 | 2.5 | 68.0 |
| PO + An.M 200 ppm | 8.0 | 4.0 | 16.0 | 2.8 | 69.0 |
| PO + An.M 600 ppm | 6.5 | 3.2 | 14.4 | 2.2 | 70.0 |
| PO + An.M 1000 ppm | 6.0 | 2.8 | 12.8 | 1.8 | 71.5 |
| PO + An.M 1400 ppm | 5.5 | 2.5 | 11.5 | 1.5 | 73.0 |
| PO + An.M 1800 ppm | 5.0 | 2.2 | 10.4 | 1.3 | 74.5 |

p-Anisidine Value

Production of secondary oxidation products, as indicated by p-anisidine value, was much lower ($P < 0.001$) in oils containing soursop flower extract than the control, which was another indication that the extract was effective in inhibiting the production of secondary oxidation products. This action had the same effect as in the case of BHT and showed that the antioxidants present in the extract are phenolic in nature and inhibit the breakdown of hydroperoxides.

TOTOX Value

TOTOX value that quantifies both hydroperoxides and products of hydroxyl radicals was less in oils with soursop flower extract than in control, indicating the potential of the extract in decreasing total oxidation. TOTOX values of oils containing soursop extract were concentration-dependent whereby the high concentrations of the extracts offered a better oxidative stability.

Thiobarbituric acid (TBA) Value

The TBA test revealed that the oils with the soursop flower extract had lower TBA values compared to the control, which is a low amount of malonaldehyde formation, a secondary oxidation product. The TBA value of the oils containing the concentration of 600-1800 ppm

was also found to be much lower ($P < 0.001$) in as compared to the control and this further substantiates the antioxidant properties of the extract.

Iodine Value

The values of iodine which show the unsaturated fatty acids were reduced with the course of storage. Nonetheless, oils with the addition of soursop flower extract experienced much less reduction of the iodine value than the control which means that the extract was effective in preventing the oxidation of unsaturated fatty acids. The extract activity was similar to that of BHT.

Soursop Flower Extract on Linoleic Acid Profile of Palm Olein

The effect of soursop flower extract on a palm olein in terms of linoleic acid profile during storage. Palm olein had a level of 10.97 of linoleic acid. There was a reduction in the level of linoleic acid content of all the samples after 30 days of the heating process where oils with the extract retained more linoleic acid than the control and the BHT-treated samples. This implies that the extract was effective in preventing oxidative degradation of linoleic acid, which again justified its possible role as a natural antioxidant.



CONCLUSION

To conclude, the *Annonamuricata* (soursop) flower extract is a potential natural antioxidant in stabilizing palm olein in the accelerated oxidation. The extract contributed greatly to oxidative stability of palm

olein, and improved the results than BHT in some tests. Also, the maintenance of the level of linoleic acid and prevention of primary and secondary oxidation products prove the usefulness of the extract as a natural substitute of synthetic antioxidants in edible oils.

REFERENCES

1. Cuvelier, M. E., & Maillard, M. N. (2012). Stabilité des huiles alimentaires au cours de leur stockage. *Oléagineux Corps Gras Lipides*, 19, 125–132.
2. Iqbal, S., & Bhanger, M. I. (2007). Stabilization of sunflower oil by garlic extract during accelerated storage. *Food Chemistry*, 100, 246–254.
3. Iqbal, S., Haleem, S., Akhtar, M., Zia-ul-Haq, M., & Akbar, J. (2008). Efficiency of pomegranate peel extracts in stabilization of sunflower oil under accelerated conditions. *Food Research International*, 45, 194–200.
4. Chang, S. S., Ostric-Matijasevic, B., Hsieh, O. A. L., & Cheng, L. H. (1977). Natural antioxidants from rosemary and sage. *Journal of Food Science*, 42, 1102–1106.
5. Thorat, I. D., Jagtap, D. D., Mohapatia, D., Joshi, D. C., Sutar, R. F., & Kapdi, S. S. (2013). Antioxidants, their properties, uses in food products and their legal implications. *International Journal of Food Studies*, 2, 81–104.
6. Krishnaiah, D., Sarbatly, R., & Nithyanandam, R. (2010). A review of the antioxidant potential of medicinal plant species. *Food Bioproducts Processing*, 157, 1–17.
7. Womeni, H. M., Tonfack, D. F., Tiencheu, B., & Linder, M. (2013). Antioxidant potential of methanolic extracts and powders of some Cameroonian spices during accelerated storage of soybean oil. *Advances in Biological Chemistry*, 3, 304–313.
8. Baskar, R., Rajeswari, V., & Kumar, T. S. (2007). In vitro antioxidant studies in leaves of *Annona* species. *Indian Journal of Experimental Biology*, 45, 480–485.
9. Olakunle, S., Onyechi, O., & James, O. (2014). Toxicity, anti-lipid peroxidation, in vitro and in vivo evaluation of antioxidant activity of *Annona muricata* ethanol stem bark extract. *American Journal of Life Sciences*, 2, 271–277.
10. Gao, X., Ohlander, M., Jeppsson, N., Björk, L., & Trajkovski, V. (2000). Changes in antioxidant effects and their relationship to phytonutrients in fruits of sea buckthorn (*Hippophae rhamnoides* L.) during maturation. *Journal of Agricultural and Food Chemistry*, 48, 1485–1490.
11. Braca, A., Sortino, C., Politi, M., Morelli, I., & Mendez, J. (2002). Antioxidant activity of flavonoids from *Licania licaniaeflora*. *Journal of Ethnopharmacology*, 79, 379–381.
12. International Dairy Federation. (1991). IDF-Square Vergote 41, Brussels, Belgium, sec. 74A.
13. Shahidi, F., & Wanasundara, U. N. (2008). Methods for measuring oxidative stability in edible oils. In C. C. Akoh & D. B. Min (Eds.), *Food lipids: Chemistry, nutrition and biotechnology* (pp. 387–388). CRC Press.
14. Draper, H. H., & Hadley, M. (1990). Malondialdehyde determination as index of lipid peroxidation. *Methods in Enzymology*, 186, 421–431.

