



CLINICAL STUDY ASSOCIATED WITH RISK FACTORS OF CANCER OF THE UTERINE CERVIX AMONG REPRODUCTIVE FEMALES IN INDIA


Pavithra P*

Assistant Professor, Department of Pathology, Kasturba Medical College, Mangaluru, India

ABSTRACT

Despite the truth that cervical most cancers is preventable disorder, it is the fourth most often identified most cancers and main purpose of cancer dying in women. Awareness regarding cervical most cancers and its prevention is pretty low amongst Indian women. The Pap test is a easy and cost powerful method for early prognosis of cervical cancer. It is necessary to make reproductive females aware of cervical cancer, so that they can impart knowledge regarding cervical cancer and its prevention to the general public. Therefore, the aim of this study was to find out the Prevalence and clinical study associated with risk factors of cancer of the uterine cervix among reproductive females. The present study was conducted in the Department of Gynecology and Obstetrics and pathology, Kasturba Medical College Mangaluru, from Oct 2016 to Oct 2017. Each woman under investigation had a clinical history, gynecological examination and pap smear. A cervical biopsy turned into also taken for histopathology. The frequency of cervical cancer was found to be about 75% in the present study. In conclusion, having multiple sexual partners, being Human Immune Deficiency Virus (HIV) fine, having STI history and early initiation of sexual intercourse had been factors associated with cervical cancer. Therefore, warding off more than one sexual companion, delaying of early sexual contact, and self-safety from STI infections may help to save you cervical cancer.

Keywords :- Human Papillomavirus, Squamocolumnar junction, Visual Inspection with Acetic Acid, Cervical Cancer.

Access this article online		
Home page: www.mcmed.us/journal/abs	Quick Response code 	
Received:25.12.17	Revised:04.01.18	Accepted:19.01.18

INTRODUCTION

Cervical cancers is the second maximum frequent malignant tumor in reproductive females worldwide, which seriously threatens women' fitness. Constant contamination of high-danger human papillomavirus (HPV) has been clarified because the required reason of cervical cancers. The clean etiology expanded the status quo and implementation of comprehensive avoidance and control device of cervical cancer.¹ In May 2018, the World Health Organization (WHO) issued a call to dispose of cervical cancer globally, and greater than 70 nations and worldwide academic societies acted definitely at once.²

Subsequently, in November 17, 2020, WHO launched the global approach to accelerate the removal of cervical cancer as a public health problem to light the street of cervical cancer deterrence and control in the destiny which suggest that 194 international locations promise together to put off cervical most cancers for the primary time³.

It is a deadly disease as soon as it reaches the invasive tiers, however out of all of the woman genital tract cancers; its miles the most effective avertable most cancers if detected at its early ranges.

Corresponding Author: Pavithra P Email: drpebyreddy@gmail.com

Population-based totally screening with Pap smear is a critical secondary safety measure for cervical most cancers that results in a excessive-treatment price among cervical most cancers sufferers. The centers to perform Pap smear are to be had inside the institute in which the examiner has been carried out Also, under the National Cancer Control Programme, screening camps for early detection of cervical cancer are organized in numerous regions of Gujarat at regular durations by using the Gujarat Cancer Research Institute which is one of the local most cancers care institutes of India.

Epidemiological research have shown the high-hazard Human Papillomavirus (HPV) to be the maximum critical risk thing and are found in ninety nine.7% of the invasive cervical most cancers worldwide. Young age, early marriage, multiple sexual companions, terrible genital hygiene, records of abortions, high parity, tobacco and oral contraceptive use, cigarette smoking, developing CaCx.⁴ It is essential to make the reproductive female aware of cervical cancer, impart knowledge regarding cervical most cancers and its prevention to the overall public. The present study changed into achieved Prevalence and clinical study associated with risk factors of cancer of the uterine cervix among reproductive females.

MATERIAL METHODS

The present study was conducted in the Department of Gynecology and Obstetrics and pathology, Kasturba Medical College Mangaluru, from Oct 2016 to Oct 2017. Females were having complains of excessive vaginal discharge, post-coital bleeding, post-menopausal bleeding, history of pain in lower abdomen etc. On examination of speculum suspected cervical lesion and unhealthy cervix. The patients underwent gynecological examination & biopsy was taken. The histological study of the samples as well as pap smear

test was performed by the Department of Pathology at Kasturba Medical College Mangaluru. A questionnaire was used to obtain information on occupation, past medical history, smoking, sexual and reproductive factors including age at marriage, age at first issue and parity. A total 367 females were screener clinic used visual inspection with acetic acid (VIA) methods. In this procedure cervix is visualized with the naked eye under a good light source at least 1 minute after applying 3% to 5% acetic acid. If the test results are negative, no aceto white lesions will be found. If the VIA test result is positive, it shows Sharp, distinct, well-defined, dense opaque acetowhite area with or without raised margins touching the squamocolumnar junction (SCJ); leukoplakia and warts. Data analysis was done using statistical package for social sciences software. Data is summarized in form of tables. Univariate logistic regression analysis was used to evaluate the factors significantly associated with cervical cancer. Multiple Logistic Regression was used to calculate the adjusted Odds ratio with 95% CI.

RESULTS

Out of 240 reproductive females, 44% belonged to the age group of 41–50 years. The average mean was 47 years of the study population. The majority of respondents (76%) were married (Table 1). 75% of appellant had some knowledge of cervical carcinoma. As per information regarding knowledge of the symptoms of cervical cancer, 169(94%) respondents stated vaginal discharge as one of the major symptoms. The percentages of respondents who mentioned menstrual abnormality and pain as symptoms were 86.9 and 66.6, respectively. Ninety six (53%) appellant, were aware of multiple sexual partners as one of the risk factors of cervical carcinoma.

Table 1 Socio-demographic profile of study population

Socio-demographic variable	Frequency	Percentage
Age		
31-40	59	24.5%
41-50	105	44.5%
51-60	69	28.7%
61-70	4	2%
71-80	3	1.5%
Marital status		
Married	183	76.2%
non married	57	23.7%
widow		
Socio-economic class		
Class1	196	81.6%
Class 2	44	18.3%

Out of 180 appellants who had some knowledge regarding cervical carcinoma, 120(67%) had knowledge regarding Pap test as one of the preventive measures (Table 2). 53.3% was found to be VIA positive. From

those who were screened positive with VIA screening test, about 69.7% were identified with high grade lesions (grade II and III) while 30.7% of them were found with low-grade lesions (Grade I).

Table 2 Comprehension regarding various aspects of carcinoma cervix

Knowledge about carcinoma cervix	Frequency	Percentage	P value
Symptoms			
Menstrual abnormality	152	84%	≥ 0.05
Vaginal discharge	169	94%	-
Pains	146	81%	< 0.05
Others(post-coital bleeding, bladder and rectal involvement, weight loss and loss of appetite)	56	31%	< 0.05
Risk factors			
Early marriage	52	29%	< 0.05
Early pregnancy	169	94%	< 0.05
Repeated pregnancy	158	88%	≥ 0.05
Oral contraceptives	132	73%	< 0.05
Multiple sexual partners	96	53%	< 0.05
Preventive measures			
Good genital hygiene	119	66.1%	< 0.05
Use of condom	156	86.6%	< 0.05
PAP test	164	91%	< 0.05
Total	180	100	

Table 3 : Knowledge about PAP TEST as screening test for carcinoma cervix.

Knowledge about PAP test as screening test for carcinoma cervix	Under went PAP test	Total
Present	120	67%
absent	60	25%
Total	240	100%

Table 4 Lesion distributions by age category among women attended cervical cancer screening center.

VIA (visual inspection of cervix with acetic acid.)-screening result		Lesion grade	
Positive	Negative	Low grade-I	High grade-II
96	84	29	67
53.3%	46.6%	30.2%	69.7%

DISCUSSION

This study was conducted along with reproductive females in order to estimate their awareness about cervical carcinoma. 75% of reproductive females had some understanding linked to cancer of the cervix. In our study, 94.2% and 84% mentioned abnormal vaginal discharge and menstrual abnormality respectively, as symptoms of cervical cancer, which is correlated with Nganwai *et al*⁵ study. Anya *et al*⁶ was found in a study among menstrual abnormality –80.6% which include post-coital bleeding, inter-menstrual bleeding and abnormal leucorrhoea or blood-stained vaginal discharge. A similar finding to our study (84%).

In present study, showed that only 53% mentioned multiple sexual partners as one of the risk

factors, asimilar findings Ali *et al*⁷, and McCarey *et al*⁸ 45%, 41% respectively mentioned multiple partners and other promiscuous behaviour as the most common risk factor. In the present study, 94% mentioned early age at pregnancy as one of the risk factors for cervical cancer. Which is correlated with Nganwai *et al* in their study mentioned that 81.8 and 85.6% of respondents knew that first sexual intercourse at a young age and having multiple sexual partners were risk factors for cervical cancer.

In our study, knowledge regarding Pap test was present in 67% of respondents. Which are correlated findings (83%) were documented in a study carried out by Mutyaba *et al*⁹

CONCLUSION

The incidence of cervical cancer was high in present study. Age at first sexual intercourse, having multiple sexual partners, being HIV positive, and having history of STI status were the known as factors related

with cervical cancer. Therefore, health programmers should be initiated to raise awareness to increase the informed health seeking behavior in the high-risk group of reproductive females.

REFERENCES

1. Garland SM, Giuliano A, Brotherton J. (2018). IPVS statement moving towards elimination of cervical cancer as a public health problem. *Papillomavirus Res*, 5:87-8.
2. Herrero R. (2018). Elimination of cervical cancer in Latin America. *Salud Publica Mex* (in Spanish), 60:621-3.
3. A cervical cancer-free future: First-ever global commitment to eliminate a cancer. Available online: <https://www.paho.org/en/news/17-11-2020-cervicalcancer-free-future-first-ever-global-commitmenteliminate-cancer>.
4. Misra JS, Srivastava S, Singh U, Srivastava AN (2009). Risk factors and strategies for control of carcinoma cervix in India: Hospital based cytological screening experience of 35 years. *Indian Journal of Cancer*, 46: 155-159.
5. Nganwai P, Truadpon P, Inpa C, Sangpetngam B, Mekjarasnapa M. (2008) Knowledge, attitudes and practices vis-a-vis cervical cancer among registered nurses at the Faculty of Medicine, Khon Kaen University, Thailand *Asian Pac J Cancer Prev*, 9:15–18.
6. Anya SE, Oshi DC, Nwosu SO. (2005) Knowledge, attitude and practice of female health professionals regarding cervical cancer and Pap smear *Niger J Med*, 14 283–6
7. Ali SF, Ayub S, Manzoor NF, Azim S, Afif M. (2010) Knowledge and awareness about cervical cancer and its prevention amongst interns and nursing staff in tertiary care hospitals in Karachi, Pakistan *PLoS ONE*, 5(6) e11059.
8. McCarey C. (2011) Awareness of HPV and cervical cancer prevention among Cameroonian healthcare workers. *BMC Women's Health* 11 45. <http://www.biomedcentral.com/1472-6874/11/45>.
9. Mutyaba TF, Mmiro A. (2006) Knowledge, attitudes and practices on cervical cancer screening among the medical workers of Mulago Hospital, Uganda *BMC Med Educ*, 6 13.

Cite this article:

Pavithra P (2018). Clinical Study Associated with Risk Factors of Cancer of the Uterine Cervix among Reproductive Females in India. *Acta Biomedica Scientia*, 5(2), 194-197



Attribution-NonCommercial-NoDerivatives 4.0 International