



## TO STUDY CONGENITAL MIDLINE NECK SWELLINGS IN SOUTH INDIAN CHILDREN'S

Venkata Sivakrishna K<sup>1</sup>, Viswateja Chitturi<sup>2\*</sup>

<sup>1</sup>Associate Professor of Cardiology, Sree Balaji Medical College and Hospital, Chennai. (Affiliated to Bharath University, Chennai) India.

<sup>2</sup>Assistant Professor of Paediatrics, Sri Lakshmi Narayana Institute of Medical Sciences, Pondicherry. (Affiliated to Bharath University, Chennai, India.

### ABSTRACT

Midline congenital neck mass is not a rare entity in children presenting to Otorhinolaryngology and Paediatric surgery department, but nothing significant is mentioned in literature, especially in our country. This study is an attempt to highlight the incidence of midline congenital neck masses and importance of early diagnosis and timely medical or surgical intervention. Cross sectional retrospective study conducted at Department of Paediatric and Cardiology, Sree Balaji Medical College and Hospital, Chennai. Data of all children below the age of 18 years presenting with congenital neck mass to our institute retrospectively analysed with regards to age, gender, types of swelling, location and the investigations performed. Data was presented in the form of tables, diagrams and pie charts. 67 patients with different congenital neck swellings were studied. Out of these most frequent were thyroglossal duct anomalies 37 (55.2%) followed by dermoid cysts 16 (23.9%), lymphangiomas 7 (10.45%) and haemangiomas 4 (5.9%). Other rare swellings included cervical teratoma, ectopic thyroid and giant congenital hamartoma with one case each. Congenital midline neck masses constitute less talked about but important clinical entity. These masses pose diagnostic, therapeutic and surgical challenges to many clinicians. Hence, for proper management, sound knowledge of these masses and adequate surgical skills are necessary prerequisites..

**Keywords :-** Congenital neck mass, Thyroglossal cyst, Dermoid cyst.

Access this article online

Home Page:  
[www.mcmed.us/journal/abs](http://www.mcmed.us/journal/abs)

Quick Response code



Received: 21.01.2023

Revised: 12.02.2023

Accepted: 18.02.2023

### INTRODUCTION

Neck masses in children is a common presentation in paediatric surgical centres [1, 2]. These are usually either congenital/developmental, inflammatory/reactive or neoplastic [3]. Congenital neck masses commonly include thyroglossal cysts, branchial cleft anomalies, dermoid cysts and vascular malformations. Other rare lesions are cervical teratomas, ectopic thyroid, midline cervical clefts [4]. Thyroglossal duct cyst is the most common midline neck mass accounting for 70% of the congenital neck anomalies [5, 6]. Correct diagnosis, and complete surgical excision require accurate knowledge of these neck masses.

Objective of this study was to determine frequency of various types of congenital midline neck swellings, their age & gender distribution and outcome of surgical management in terms of recurrence.

### MATERIAL AND METHODS

A retrospective study was conducted to analyze midline congenital neck masses in paediatric age group in our institute. Records of patients for detailed history and examination were retrieved from Department of paediatric and Cardiology Department, Sree Balaji Medical College and Hospital, and were analyzed and statistical analysis was done

Corresponding Author **Dr. Viswateja Chitturi** Email: drnvsreddy2000@gmail.com

## RESULTS:

Within 10 year period, total 67 cases of midline neck masses were found. Out of these, most frequent were thyroglossal duct anomalies 37 (55.2%) followed by dermoid cysts 16 (23.9%), haemangiomas 4 (5.9%) and lymphangiomas 7 (10.45%). Other rare swellings included cervical teratoma, ectopic thyroid and giant congenital hamartoma with one case each (Fig. 1).

Age of presentation varied from 4 days to 15 yrs, majority of them lying in 4 to 6 years group closely followed by 0 to 2 yrs age group. There was a male preponderance with 42 males and 25 females giving M:F ratio of 1.68:1 (Fig.No. 2).

Out of 37 cases of thyroglossal duct anomalies, 19 cases (51.35%) were of thyroglossal fistula (Fig.4a) and 18 cases (48.65%) were of thyroglossal cyst (Fig.4b). All cases of thyroglossal fistula were iatrogenic in origin i.e. due to inadvertent incision and drainage of thyroglossal cyst abscess. Thyroglossal duct anomalies were more common in males with 25 cases (67.6%) as compared to 12 female cases (32.4%) with sex ratio of 2.1:1. Predominant position was infrahyoid (59.4%) in 22 cases followed by suprahyoid (29.7%) in 11 cases, juxtahyoid (8.1%) in 3 cases and suprasternal (2.7%) in 1 case. Mean age of the patients with thyroglossal duct anomalies was 7.15 years (Table.No. 1).

In present series, 16 cases of dermoid cyst were reported (Fig 4c). Majority were in age group of 4 to 6 years with equivocal sex distribution with 8 cases each (Table 2).

Congenital midline vascular swelling also was not a rare entity in our study. 4 cases of haemangioma (Fig.3d)

were reported with 2 cases each in 0-2 year and 2 -4 year age group with no gender predilection. Major presenting complaint was ulceration and bleeding in 75% cases. 7 cases of lymphangioma were reported out of which 2 cases showed cervico-mediastinal extension. Cysts were infected in 3 cases (Fig.4a) and non-infected (Fig.4b) in remaining 4 cases. Sex ratio was 2.5:1 and presenting age group was 0-2 years in all cases.

One case each of ectopic thyroid (8 yr male), cervical teratoma (4 day female) (Fig.4c) and giant congenital hamartoma (Fig.4d) was also reported.

Diagnosis was mainly clinical though ultrasonography and Fine Needle Aspiration Cytology(FNAC) was done to confirm the diagnosis. FNAC was particularly relevant to rule out ectopic thyroid, as misdiagnosed excision can lead to permanent hypothyroidism. CT scan provides additional information regarding extent and inner composition of mass. Chest X-ray is important in cases of cystic hygroma to rule out the mediastinal extension. All the patients diagnosed with Thyroglossal cyst/fistula underwent Sistrunk procedure under general anaesthesia. In this procedure, cyst was removed along with central portion of hyoid bone with meticulous excision of persistent thyroid duct upto foramen caecum. In patients of dermoid cysts, haemangioma, lymphangioma and others, simple surgical excision was done.

Regular follow up was done monthly for 1 year for any recurrence or fistula formation. 3 cases of recurrent Thyroglossal fistula were recorded who were re-operated after 1 year.


**Table 1: Age distribution of Thyroglossal duct anomalies:**

Age group (in years)	No of cases
0-2	2
2-4	6
4-6	6
6-8	5
8-10	7
10-12	6
12-14	4
14-16	1


**Table-2 Age distribution of Dermoid cysts.**

Age group(in years)	No of cases
0-2	0
2-4	3
4-6	7
6-8	2
8-10	2
10-12	2
12-14	0

**Figure 1: Different types of congenital midline neck swellings. The above table shows thyroglossal cyst to be commonest swelling (55.2%) followed by dermoid cyst (23.9%).**



**Figure 2: Sex distribution of patients with congenital neck swellings**



**Figure 3: a) thyroglossal fistula, b) thyroglossal cyst, c) dermoid, d) haemangioma**



**Figure 4: a) midline cystic hygroma (infected), b) midline cystic hygroma (non-infected), c) cervical teratoma, d) congenital giant hamartoma**



**DISCUSSION:**

Congenital neck masses are the most common non-inflammatory neck swellings. Nothing significant is documented in the literature till date regarding midline congenital neck swellings. Thus we compared our results with the data on congenital neck swellings in toto. Siddique et al studied 36 patients and showed thyroglossal cyst (58.33%) to be the commonest midline neck swelling followed by branchial cysts (19.44%), haemangiomas (8.33%), lymphangiomas (8.33%) and dermoid cysts (5.55%) [7]. Another study by Al Khateeb et al on 252 cases also revealed thyroglossal cyst (53%) as the most frequent swelling followed by branchial

cysts (22%), dermoid cysts (11%), haemangiomas (7%) and lymphangiomas (6%) [8]. Results of our study were found to be consistent with previous studies in terms of thyroglossal cyst. However, second most common midline swelling was dermoid cyst in our study in contrast to branchial cyst which was second most common swelling in all other studies [7-9].

Patients varied in age from 4 days to 15 yrs, majority of them lying in 4 to 6 years group in our study. Siddique et al [7] reported this to be 6 to 8 years while Ayugi et al [10] found it to be 0-2 years. Our study showed a male preponderance with M:F ratio of 1.58:1 which is in agreement with Siddique et al ratio of

1.11:1. However Al Khateeb et al [8] found out a female predilection with a ratio of 1:1.2 while Ayugi et al [10] had an equivocal distribution (M:F=1:1).

Thyroglossal cyst represents most common congenital midline neck swelling. It arises from persistent thyroglossal duct. Thyroglossal fistulae on the other hand are acquired following rupture or incision of infected thyroglossal cyst. Thyroglossal duct anomalies were more common in males with 25 cases (67.6%) as compared to 12 female cases (32.4%) with sex ratio of 2.1:1. In other published series, sex ratio was 1.1:1 [7] and 1:1 [11, 12].

Mean age of the patients with thyroglossal duct anomalies was 7.15 years as compared to 7.8 years, 37.6 years and 15.4 years in studies by Ayugi et al [10], Tarcoveanu E et al [13] and Al Salem et al [14], respectively. Predominant position was infrahyoid (59.4%) in 22 cases followed by suprahyoid (29.7%) in 11 cases, juxtahyoid (8.1%) in 3 cases and suprasternal (2.7%) in 1 case. These findings were similar to Siddique et al [7] and Shih-Tsang L et al [12]. Thyroglossal cyst frequently gets infected leading to iatrogenic fistula formation. Thus in authors opinion, thyroglossal cyst should be excised as early as possible. Dermoid cysts are slow growing benign tumors which may occur in midline of the neck. In the neck, dermoid cysts are usually firm lumps attached to the overlying skin. In present series, 16 cases of dermoid cyst were reported. Majority were in age group of 4 to 6 years. This is contrary to Shih-Tsang et al [12] study which states that dermoid usually presents in second and third decade. There was no gender predilection.

## REFERENCES

1. Tuskington, J. R., Paterson, A., Sweeney, L. E., & Thornbury, G. D. (2005). Neck masses in children. *British Journal of Radiology*, 78, 75-85.
2. Tracy, T. F. Jr., & Muratore, C. S. (2007). Management of common head and neck masses. *Seminars in Pediatric Surgery*, 16, 3-.
3. Meier, J. D., & Grimmer, J. F. (2014). Evaluation and management of neck masses in children. *American Family Physician*, 89, 353-358.
4. LaRiviere, C. A., & Waldhausen, J. H. (2012). Congenital cervical cysts, sinuses, and fistulae in pediatric surgery. *Surgical Clinics of North America*, 92, 583-597.
5. Mondin, V., Ferlito, A., Muzzi, E., Silver, C. E., Fagan, J. J., Devaney, K. O., & Rinaldo, A. (2008). Thyroglossal duct cyst: Personal experience and literature review. *Auris Nasus Larynx*, 35, 11-25.
6. Ruchira, M., Vacha, S., Kamlesh, J., & Inderraj, T. (2010). Thyroglossal cyst: An unusual appearance. *Bombay Hospital Journal*, 52, 93-96.
7. Siddique, M. A., Hossen, M., Taous, A., Islam, S. M., & Khan, S. A. (2012). Clinical presentation of congenital neck mass in children. *Bangladesh Journal of Otorhinolaryngology*, 18, 16-22.
8. Al-Khateeb, T. H., & Zoubi, F. (2007). Congenital neck masses: A descriptive retrospective study of 252 cases. *Journal of Oral and Maxillofacial Surgery*, 65, 2242-2247.
9. Stephanie, P. A., & John, H. T. W. (2007). Congenital cervical cysts, sinuses, and fistulae. *Otolaryngologic Clinics of North America*, 40, 161-176.
10. Ayugi, J. W., Ogeng'o, J. A., & Macharia, I. M. (2010). Pattern of congenital neck masses in a Kenyan pediatric population. *International Journal of Pediatric Otorhinolaryngology*, 74, 64-66.

Congenital midline vascular swelling also was not a rare entity in our study. Haemangioma being the commonest benign tumor of infancy, affects head and neck in 14 – 20% of cases. Lymphangiomas are degenerative lesions arising from lymphatics and can be classified as simple lymphangiomas, cavernous lymphangiomas and cystic hygromas. A cystic hygroma is the most common form of lymphangioma and constitutes about 5% of all benign tumors of infancy and childhood and is thought to arise from an early sequestration of embryonic lymphatic channels [11]. All cases were in age group of 0 to 4 years equally presenting in both the sexes. Siddique et al showed that 66.67% cases reported at time of birth and the remaining in second decade with M:F= 2:1 [7]. According to Krol et al, infantile haemangioma typically appears in first few weeks of life and proliferates for weeks to several months [15]. In our opinion haemangioma of neck should also be excised as early as possible since all these cases present with bleeding and ulceration

## CONCLUSION

Congenital midline neck masses constitute less talked about but important clinical entity. These masses pose diagnostic, therapeutic and surgical challenges to many clinicians. Hence, sound knowledge of these masses, timely intervention and adequate surgical skills will not only ensure proper management but also improve the final outcome as well as prognosis. This will go a long way in improving the expectancy and quality of life in these cases.

11. Watkinson, J. C., Gaze, M. N., & Wilson, J. A. (2000). Benign neck disease. In: Stell and Maran's head and neck surgery. *Butterworth and Heinemann, Oxford*, 4, 181-186.
12. Shih-Tsang, L., Fen-Yu, T., Chuan-Jan, H., Tsai, C., & Lee, S. (2008). Thyroglossal duct cyst: A comparison between children and adults. *American Journal of Otolaryngology-Head and Neck Surgery*, 29, 83-87.
13. Tarcoveanu, E., Niculescu, D., Elena, C., Goleanu, V., & Vasilescu, A. (2009). Thyroglossal duct cyst. *Journal of Chirurgie Lasi*, 5, 75-78.
14. Al-Salem, A. H., Quasaruddin, S., & Ahmed, M. (1996). Thyroglossal cyst: A clinicopathological study. *Saudi Medical Journal*, 17, 620-625.
15. Krol, A., & MacArthur, C. J. (2005). Congenital hemangiomas: Rapidly involving and non-involving congenital hemangioma. *Archives of Facial Plastic Surgery*, 7, 307-311.

**Cite this article:**

Venkata Sivakrishna K, Viswateja Chitturi. (2023). "To Study Congenital Midline Neck Swellings in South Indian Children's". *Acta Biomedica Scientia*, 10(2): 32-36.



**Attribution-NonCommercial-NoDerivatives 4.0 International**