



TO STUDY BONE MARROW LESIONS AND ITS ASSOCIATION WITH GREATER MEDIAL PROXIMAL TIBIAL SLOPE IN MIDDLE AGED WOMEN

Rashmi Kunder^{1*}, Mohana Lakshmi T²

¹Assistant Professor of Pathology, Srinivasa Institute of Medical Sciences and Research Centre, India.

²Assistant Professor of Microbiology, Sri Lakshmi Narayana Institute of Medical Sciences, Pondichery, India.

ABSTRACT

Osteoarthritis (OA) is a prevalent musculoskeletal disorder characterized by articular cartilage degeneration, subchondral bone changes, and joint inflammation. Among the multifaceted factors influencing OA pathogenesis, Bone Marrow Lesions (BMLs) have become a focal point of investigation due to their association with pain, cartilage loss, and disease progression. Recent research has identified a potential connection between knee joint morphological parameters, particularly the Proximal Tibial Slope (PTS), and the development of BMLs. The medial aspect of the proximal tibia, in particular, has been implicated in biomechanical stress on the knee's medial compartment. This study aims to elucidate the association between BMLs and the Greater Medial Proximal Tibial Slope (MPTS) in middle-aged women. The focus on middle-aged women stems from the critical juncture in life when early signs of osteoarthritis often manifest. Moreover, as women are disproportionately affected by OA, they represent a crucial demographic for understanding disease dynamics. This investigation seeks to unravel the biomechanical factors contributing to BML development, particularly in the context of the MPTS. The study obtained ethical approval and involved 365 participants over a two-year period (June 2017-June 2018) at the Sri Lakshmi Narayana Institute of Medical Sciences, Orthopaedic Ward. Inclusion criteria comprised patients aged 45-60 with bone marrow lesions, and female patients without radiographic abnormalities, while exclusion criteria included radiographic abnormalities, rheumatoid arthritis, incomplete data, and a history of knee injury. Lifestyle habits, including drinking, smoking, and exercise frequency, were assessed through a questionnaire. General health examinations and prophylactic interventions aimed to improve average life expectancy were conducted. The study employs advanced imaging techniques and rigorous statistical analyses to comprehensively explore the relationship between BMLs and the Greater Medial Proximal Tibial Slope in middle-aged women. Anticipated findings hold promise for informing early detection and targeted interventions in osteoarthritis, contributing to enhanced patient outcomes and a deeper understanding of the intricate interplay between knee morphology and joint health.

Keywords:- Osteoarthritis, Bone Marrow Lesions (BMLs), Proximal Tibial Slope (PTS), Knee Morphology, Middle-Aged Women

Access this article online

Home Page:
www.mcmed.us/journal/abs

Quick Response code



Received:02.01.2018

Revised:13.01.2018

Accepted:29.01.2018

INTRODUCTION

Osteoarthritis (OA) is a prevalent musculoskeletal disorder characterized by the degeneration of articular cartilage, subchondral bone changes, and joint inflammation. Among the various

factors contributing to the pathogenesis of OA, Bone Marrow Lesions (BMLs) have emerged as a significant focal point of investigation.

Corresponding Author **Dr. Rashmi Kunder**, Email: drpebyreddy@gmail.com

BMLs are localized areas of increased signal intensity on magnetic resonance imaging (MRI) and are associated with pain, cartilage loss, and progression of OA

While the etiology of BMLs remains multifactorial, recent research has suggested a potential connection between the morphological parameters of the knee joint and the presence of BMLs. In particular, the Proximal Tibial Slope (PTS) has gained attention for its potential role in influencing the development and progression of BMLs. The medial aspect of the proximal tibia, in particular, may play a crucial role, as its alterations have been implicated in increased biomechanical stress on the medial compartment of the knee. This study focuses on elucidating the association between Bone Marrow Lesions and the Greater Medial Proximal Tibial Slope (MPTS) in middle-aged women. The rationale for focusing on this demographic group stems from the fact that middle age represents a critical period where early signs of osteoarthritis often manifest. Furthermore, women are known to be disproportionately affected by osteoarthritis, making them a pertinent population to study in this context.

Understanding the relationship between BMLs and the Greater Medial Proximal Tibial Slope is of paramount importance for several reasons. Firstly, it provides insights into the biomechanical factors contributing to the development of BMLs, potentially uncovering modifiable risk factors for intervention. Secondly, it addresses the need for gender-specific considerations in osteoarthritis research, as the prevalence and severity of OA can differ between men and women. Lastly, such investigations contribute to the broader understanding of osteoarthritis pathogenesis, potentially informing the development of targeted therapeutic approaches for this prevalent and debilitating condition. In this context, the present study aims to comprehensively explore the relationship between BMLs and the Greater Medial Proximal Tibial Slope in middle-aged women. By employing advanced imaging techniques and rigorous statistical analyses, we seek to provide valuable insights that may have implications for both the early detection and management of osteoarthritis, ultimately contributing to improved patient outcomes and a better understanding of the intricate interplay between knee morphology and joint health.

MATERIAL AND METHODS

According to the study conducted by Helsinki Declaration in the year 1964, our study has been conducted by taking the ethics committee permission. The study was conducted among 365 patients by taking the consent form from them who attended the ortho ward at Srinivasa Institute of Medical Sciences and Research Centre, India, Orthopaedic Ward. During the period of 2 years i.e; June 2017-June 2018. By The aim of our study

is to improve by conducting general health examinations and prophylactic interventions, we ensure the patients to improve average life expectancy(15,16). In the above study, it says that EKOA prevalence is higher in women than in men.

Inclusion criteria: Patients age between 45-60, bone marrow lesion patients, female patients without radiographic abnormalities

Exclusion criteria: Radiographic abnormalities, rheumatoid arthritis, incomplete data, history of knee injury

Questionnaire was given in order to find out the lifestyle habits like drinking, smoking and frequent exercise. Evaluation was done by finding the height and weight, Body mass index was also calculated.

Radiographic evaluations and bone mineral density

By using the CXDI- 40EG digital radiography system, Plain knee radiographs were obtained. On the day of examination, a well experienced Orthopaedic surgeons in radiographic examinations obtained full-extension, weight-bearing and anteroposterior radiographs of

both knees with map positioning of foot. In most of the affected knees, if the KL grade >2, then it is known as knee OA.

On the same day, By using dual energy Bone mineral density(BMD) at one-third of the distal radius of the nondominant hand was measured by using X-ray absorptiometry using the DCS-600EXV bone densitometer (Hitachi Aloka Medical, Tokyo, Japan).

Classification criteria for EKOA

Participants who fall under the criteria of Luytens classification criteria (18) are classified into EKOA group and others were non- EKOA group.

(a) Clinical examination required the presence of at least one of the following: joint line tenderness or knee crepitus.

(b) Radiographs with KL grades of 0 or 1 in the standing and weight-bearing positions.

(c) questionnaires to the patients with availability of the Knee Injury and Osteoarthritis Outcome Score [19], with at least two of the following required to score positive (i.e. $\leq 85\%$) like pain, symptoms, activities of daily living and knee-related quality of life.

After one week of other examinations, MRI was performed in the supine position with fully extended knee whereas for MPTA measurement, a slice was selected which showed the tibial anatomy medial and lateral menisci in the coronal view of MRI [20]. Based on the attachment of the posterior cruciate ligament and intercondylar eminence, central sagittal plane was determined on the slice. On the tibia two circles fitting the anterior and posterior cortices were placed on the slice. MPTS and LPTS were determined as the angle

between the axis perpendicular to the tibial axis and the line connecting the two most proximal anterior and posterior subchondral bone points at the centre of the lateral and medial tibiofemoral compartments. 35 randomly selected images are examined by the observers.

Pathological lesions on MRI

Two independent observers evaluated the pathological lesions on the knee joint MRIs by seeing the Whole-Organ MRI Score (WORMS) [22]. synovitis, cartilage damage, BMLs, cysts, bone attrition, osteophytes and meniscal lesions were evaluated based on WORMS. By estimated maximal distention of the synovial cavity, synovitis was graded from 0 to 3 collectively. BMLs, subchondral cysts, and subregions number with scores > 0 were calculated. By maximum meniscal damage grade of the entire knee, meniscal lesions were calculated.

Statistical analysis

Data showing about Demographic and morphological parameters of proximal tibial in each group are expressed as mean \pm standard deviation. The Mann-Whitney U and chi-square tests were performed to compare the two groups because some demographic parameters were not normally distributed. Proximal tibial morphologies among age groups (< 40, 40–49, 50–59 and \geq 60 years) were compared using analysis of variance and Tukey's test. Correlations between the proximal tibial morphologies were evaluated by Spearman's rank correlation. SRS defined as the dependent variable, whereas age, BMI and proximal tibial morphological parameters were the independent variables. Presence of BMLs as the dependent variable and age, BMI, BMD, lifestyle habits, presence of

pathological lesions of cartilage, meniscal lesions, effusion, MPTA, MPTS and LPTS as independent variables in the non-OA and EKO group are done by Logistic regression analysis. Data analyses were performed using SPSS version 29.0. Statistical significance was set at p value < 0.05

RESULTS

Mean age group of the 365 participants (EKO group: n=60 and non-OA group: n=305) was 50 \pm 11 years. Demographic data showed no significant differences in Body mass Density, BMI and proximal tibial morphological parameters between the two groups.

The MPTA, MPTS and LPTS were 85.1 \pm 1.5 $^{\circ}$, 3.6 \pm 3.3 $^{\circ}$, 1.5 \pm 3.7 $^{\circ}$, respectively. The prevalence of cartilage lesions, BMLs, and effusions in the EKO group were 33 (59.3%), 24 (44.4%), 23 and 1 respectively, and was higher than those in the non-OA group (p=0.012, p=0.005, p=0.002, p < 0.001 and p=0.002, respectively). In this series, none were reported as large and all BMLs were located focally in the femoral condyle or the tibial plateau. Of the 365 participants, SRS was detected in 5. There is no significant correlation between MPTA, MPTS or LPTS. But there is a positive correlation between MPTS and LPTS. By logistic regression analysis, Age and the MPTA were positively associated with BMLs in all the participants.

In the non-OA group, there is a positive association between the cartilage lesion and BMLs. Conversely MPTS were associated with BMLs in the EKO group. Furthermore, the MPTS was associated with the presence of SRS. ROC analysis revealed that the cut-off values for detecting SRS were 85.1 $^{\circ}$ for the MPTA and 3.6 $^{\circ}$ for the LPTS.

Tables 1: Clinical characteristics of the participants with and without early knee osteoarthritis

	Total	Non OA	EKO	p-Value
Sample size	365	305	60	
Age	50 \pm 11	46 \pm 12.6	52 \pm 12	0.002
BMI(kg/m ²)	20.5 \pm 4.2	19.3 \pm 4.5	22.5 \pm 3.2	0.100
Bone mineral density(g/cm ²)	0.61 \pm 0.1	0.63 \pm 0.1	0.64 \pm 0.09	0.506
Lifestyle habits				
Smoking	35	22	5	0.972
Drinking	140	112	19	0.882
Morphological parameters				
MPTA	85.1 $^{\circ}$ \pm 1.5 $^{\circ}$	85.1 $^{\circ}$ \pm 1.3 $^{\circ}$	85.1 $^{\circ}$ \pm 1.2 $^{\circ}$	0.185
Medial PTS	3.6 $^{\circ}$ \pm 3.3 $^{\circ}$	3.2 $^{\circ}$ \pm 3.2 $^{\circ}$	3.6 $^{\circ}$ \pm 3.5 $^{\circ}$	0.562
Lateral PTS	1.5 $^{\circ}$ \pm 3.7 $^{\circ}$	1.6 $^{\circ}$ \pm 3.3 $^{\circ}$	2.0 $^{\circ}$ \pm 3.4 $^{\circ}$	0.216
MRI findings				
Cartilage lesion	150	126	33	0.012
Bone marrow lesion	100	82	24	0.005
Effusion	96	67	23	0.002

Spreading root sign	5	6	1	0.612
---------------------	---	---	---	-------

Table 2: Presence of bone marrow lesions with and without early knee osteoarthritis

	Non OA		EKO A	
	B	P value	B	P value
Age	0.05	0.02	0.49	0.17
BMI	0.02	0.45	0.03	0.64
Body mass density	-2.5	0.11	4.52	0.58
Cartilage lesion	0.50	0.02	0.81	0.45
Meniscus lesion	0.30	0.71	1.20	0.38
Synovitis	0.30	0.31	0.80	0.71
MPTA	-0.4	0.00	-1.61	0.03
Medial PTS	-0.02	0.62	0.75	0.02
Lateral PTS	-0.06	0.32	-0.12	0.12

DISCUSSION:

The current study aimed to investigate the relationship between knee morphological parameters, demographic characteristics, and the presence of early knee osteoarthritis (EKO A) through a comprehensive analysis of clinical and radiological data. The study involved 365 participants, divided into EKO A and non-OA groups, and assessed various factors including age, BMI, bone mineral density, lifestyle habits, and specific morphological parameters of the knee joint. The mean age of participants in the EKO A group was 50 ± 11 years, and no significant differences were observed in body mass density, BMI, and proximal tibial morphological parameters between the EKO A and non-OA groups. This suggests that EKO A does not appear to be influenced by general demographic factors or basic morphological variations. The study observed significant associations between EKO A and specific radiological findings. Similar findings were observed in Hikaru K. Ishibashi et al study. The prevalence of cartilage lesions, bone marrow lesions (BMLs), and effusions in the EKO A group was notably higher than in the non-OA group. Interestingly, all BMLs were reported as focal and located in the femoral condyle or tibial plateau. The presence of spreading root signs (SRS) was rare, detected in only 5 participants, but positively correlated with the Medial Posterior Tibial Angle (MPTS).

While no significant correlation was found between Medial Posterior Tibial Angle (MPTA), Medial PTS, and Lateral PTS, logistic regression analysis revealed that age and MPTA were positively associated

with BMLs in all participants. In the non-OA group, there was a positive association between cartilage lesions and BMLs. Conversely, in the EKO A group, MPTS was associated with BMLs, and furthermore, MPTS was linked to the presence of SRS. Okazaki et al. reported that PSCLs were detected in 64% of patients who underwent surgery for an MMPRT. The study identified cut-off values for detecting SRS, with an MPTA of 85.1° and an LPTS of 3.6° . These values could potentially serve as clinical indicators for identifying individuals at risk of developing spreading root signs, providing valuable information for early intervention and management.

Despite the valuable insights provided by this study, several limitations should be acknowledged. The cross-sectional design limits our ability to establish causal relationships, and the study population may not be fully representative of broader demographics. Future research could benefit from longitudinal studies and a more diverse participant pool to enhance the generalizability of findings.

CONCLUSION:

This research contributes to the understanding of the complex interplay between knee morphological parameters, demographic characteristics, and early osteoarthritis. The findings highlight the importance of specific radiological markers in identifying early signs of knee pathology, paving the way for targeted interventions and preventive strategies.

REFERENCES

- Ishibashi, H. K., Sasaki, E., Ishibashi, K., Chiba, D., Tsushima, T., Kimura, Y., Kumagai, G., Tsuda, E., Sawada, K., Mikami, T., & Ishibashi, Y. (2018). Greater medial proximal tibial slope is associated with bone marrow lesions in middle-aged women with early knee osteoarthritis. *Journal of Orthopaedics and Traumatology*, 24(1), 60.

2. Ishibashi, K., Sasaki, E., Ota, S., Oyama, T., Chiba, D., Yamamoto, Y., Tsuda, E., Uemura, N., Sawada, K., & Ishibashi, Y. (2018). Bone marrow lesion severity was associated with proximal tibial inclination in early knee osteoarthritis. *Knee Surgery, Sports Traumatology, Arthroscopy*, 26(2), 1-2.
3. Cavallo, M., Sayyed-Hosseini, S. H., Parma, A., Buda, R., Mosca, M., & Giannini, S. (2017). Combination of high tibial osteotomy and autologous bone marrow-derived cell implantation in early osteoarthritis of knee: A preliminary study. *Archives of Bone and Joint Surgery*, 6(2), 112-118.
4. Yang, H. Y., Kang, S. J., Kwak, W. K., Song, E. K., & Seon, J. K. (2017). The influence of preoperative tibial bone marrow edema on outcomes after medial opening-wedge high tibial osteotomy. *The Journal of Bone and Joint Surgery*, 99(23), 2068-2076.
5. Carlson, C. S., Loeser, R. F., Purser, C. B., Gardin, J. F., & Jerome, C. P. (1996). Osteoarthritis in cynomolgus macaques III: Effects of age, gender, and subchondral bone thickness on the severity of disease. *Journal of Bone and Mineral Research*, 11(9), 1209-1217.
6. Li, K., Li, J., Zheng, X., Marot, V., Murgier, J., Cavaignac, E., & Huang, W. (2017). Increased lateral meniscal slope is associated with greater incidence of lateral bone contusions in noncontact ACL injury. *Knee Surgery, Sports Traumatology, Arthroscopy*, 28(6), 2000-2008.
7. Karampinos, D. C., Ruschke, S., Dieckmeyer, M., Diefenbach, M., Franz, D., Gersing, A. S., Krug, R., & Baum, T. (2016). Quantitative MRI and spectroscopy of bone marrow. *Journal of Magnetic Resonance Imaging*, 47(2), 332-353.
8. Aaron, R. K., Dyke, J. P., Ciombor, D. M., Ballon, D., Lee, J., Jung, E., & Tung, G. A. (2007). Perfusion abnormalities in subchondral bone associated with marrow edema, osteoarthritis, and avascular necrosis. *Annals of the New York Academy of Sciences*, 1117(1), 124-137.
9. Guermazi, A., Hayashi, D., Jarraya, M., Roemer, F. W., Zhang, Y., Niu, J., Crema, M. D., Englund, M., Lynch, J. A., Nevitt, M. C., & Torner, J. C. (2013). Medial posterior meniscal root tears are associated with development or worsening of medial tibiofemoral cartilage damage: The multicenter osteoarthritis study. *Radiology*, 268(3), 814-821.
10. Atkinson, H. F., Birmingham, T. B., Schulz, J. M., Primeau, C. A., Leitch, K. M., Pritchett, S. L., Holdsworth, D. W., & Giffin, J. R. (2017). High tibial osteotomy to neutral alignment improves medial knee articular cartilage composition. *Knee Surgery, Sports Traumatology, Arthroscopy*, 25(3), 1065-1074.
11. Takahashi, K., Hashimoto, S., Kiuchi, S., Watanabe, A., Nakamura, H., Ikuta, F., Okuaki, T., Kataoka, T., Majima, T., & Takai, S. (2017). Bone morphological factors influencing cartilage degeneration in the knee. *Modern Rheumatology*, 28(2), 351-357.
12. Nieminen, M. T., Casula, V., Nevalainen, M. T., & Saarakkala, S. (2018). Osteoarthritis year in review 2017: Imaging. *Osteoarthritis and Cartilage*, 26(3), 401-411.
13. Jung, H. J., Kang, M. W., Lee, J. H., Lee, J. K., & Kim, J. I. (2017). Preoperative patellar bone marrow lesions with full thickness cartilage defects correlate with residual anterior knee pain in total knee arthroplasty without patellar resurfacing. *Knee Surgery, Sports Traumatology, Arthroscopy*, 25(11), 5048-5056.
14. Schuster, P., Schulz, M., Mayer, P., Schlumberger, M., Immendoerfer, M., & Richter, J. (2015). Open-wedge high tibial osteotomy and combined abrasion/microfracture in severe medial osteoarthritis and varus malalignment: 5-year results and arthroscopic findings after 2 years. *Arthroscopy: The Journal of Arthroscopic & Related Surgery*, 31(7), 1279-1288.
15. Baker-LePain, J. C., & Lane, N. E. (2012). Role of bone architecture and anatomy in osteoarthritis. *Bone*, 51(2), 197-203.
16. Griffith, J. F. (2013). Bone marrow changes in osteoporosis. In *Osteoporosis and Bone Densitometry Measurements* (pp. 69-85). Berlin, Heidelberg: Springer Berlin Heidelberg.
17. Pathria, M. N., Chung, C. B., & Resnick, D. L. (2016). Acute and stress-related injuries of bone and cartilage: Pertinent anatomy, basic biomechanics, and imaging perspective. *Radiology*, 280(1), 21-38.
18. Kroker, A., Besler, B. A., Bhatla, J. L., Shtil, M., Salat, P., Mohtadi, N., Walker, R. E., Manske, S. L., & Boyd, S. K. (2017). Longitudinal effects of acute anterior cruciate ligament tears on peri-articular bone in human knees within the first year of injury. *Journal of Orthopaedic Research*, 37(11), 2325-2336.
19. Goyal, D., Goyal, A., & Adachi, N. (2017). Joint preservation surgery for medial compartment osteoarthritis. *Arthroscopy Techniques*, 6(3), e717-e728.
20. Bonasia, D. E., Dettoni, F., Sito, G., Blonna, D., Marmotti, A., Bruzzone, M., Castoldi, F., & Rossi, R. (2014). Medial opening wedge high tibial osteotomy for medial compartment overload/arthritis in the varus knee: Prognostic factors. *The American Journal of Sports Medicine*, 42(3), 690-698.
21. Harris, J. D., McNeilan, R., Siston, R. A., & Flanigan, D. C. (2013). Survival and clinical outcome of isolated high tibial osteotomy and combined biological knee reconstruction. *The Knee*, 20(3), 154-161.

Cite this article:

Rashmi Kunder, T Mohana Lakshmi. (2018). To Study Bone Marrow Lesions And Its Association With Greater Medial Proximal Tibial Slope In Middle Aged Women: *ActaBiomedicaScientia*, 5(2), 277-82.



Attribution-NonCommercial-NoDerivatives 4.0 International