



## CANADIAN JOURNAL OF BIOLOGICAL AND MICROBIOLOGY RESEARCH



Journal homepage: [www. http://mcmed.us/journal/cjbr](http://mcmed.us/journal/cjbr)

### A STUDY ON CHEMICAL AND BACTERIOLOGICAL ANALYSIS OF BORE WATER FROM SOME COMMUNITIES IN TIRUPATHI RURAL AREAS OF CHITTOOR DISTRICT, ANDHRA PRADESH, INDIA

**N. Supraja**

Ph.D Research Scholar, Department of Biotechnology, Thiruvalluvar University, Vellore-632115, Tamilnadu, India.

#### Article Info

Received 25/08/2013

Revised 15/09/2013

Accepted 18/10/2013

**Key words:** bore water, Tirupathi Rural region areas, bacteria, Trace metals, physiochemical Properties.

#### ABSTRACT

This study consisted the determination of trace metal ions, bacteriological analyses and some physiochemical properties in bore water samples from some communities in tirupathi rural areas of Chittoor district, where bore water samples are not treated before it is consumed. The purpose was to ascertain the quality of water from these sources. Samples were taken from four sampling points and analyzed for the following parameters Fluorides, Chlorides, Calcium carbonate and Nitrites. The presence of pathogens in water for drinking purposes is of public health significance considering the possibility of the presence of other bacteria, protozoa and enteric viruses that are implicated in gastro-intestinal water borne diseases and the low infectious dose for these water borne pathogens.

#### INTRODUCTION

Good drinking water quality is essential for the well-being of all people. Unfortunately in many countries around the world, including India, some drinking water supplies have become contaminated, which has impacted on the health and economic status of the populations. Contaminants such as bacteria, viruses, nitrates and salt have found their way into water supplies as a result of inadequate treatment and disposal of waste industrial discharges, and over-use of limited water resources. Even other Chemicals to be harmful to human health. Unfortunately, this problem arose because the bore water was extracted for drinking without a detailed chemical investigation. The natural water analyses for physical and chemical properties including trace element contents are very important for public health studies. The bacterial qualities of bore water, pipe borne water and other natural water supplies in tirupathi, have been reported to be unsatisfactory, with coli form counts far exceeding the

level recommendation by W.H.O [1-3]. The reason for elucidation of important parameters in water quality assessment may be attributed to the fact that in the overall potability of water, such parameters should not be ignored [4]. According to our literature review some physical and chemical properties of the samples were determined by using standard analytical methods. The probability of ingesting infective dose of disease causing microorganism is very high considering the fact that water borne pathogens generally have low infective dose. The objective of this work is to evaluate the general bacteriological and physico chemical parameters of the sources of water used for drinking purposes from bore.

#### MATERIALS AND METHODS

##### Sample collection

The bore water samples were collected in prewashed (with detergent, diluted HNO<sub>3</sub> and doubly deionized distilled water, respectively) polyethylene bottles. pH and conductivity of the samples were measured while collecting the samples. Water samples from four different areas located in and around tirupathi territory were collected in brown glass bottles with necessary precautions. Samples were collected in July 2013. The determinations of the physicochemical and Bacteriological

Corresponding Author

**N. Supraja**

Email:- [krishna.supraja@gmail.com](mailto:krishna.supraja@gmail.com)



properties of the water samples were performed on the same day of sampling. The water samples were obtained directly from the bore after allowing the water to run for at least five minutes and each sample bottle and its cap rinsed three times. These samples were subsequently stored at 4°C for as short a time as possible before analysis to minimize physicochemical changes [5]. Because very little particulate matter was present in the sample, filtration was not considered necessary.

### Physicochemical Analysis

The physicochemical tests included the determination of temperature, turbidity, odour, colour, pH, conductivity, nitrites content, alkalinity, total hardness, fluorides and chloride content using the methods of FAO (1997a) [6].

### Bacteriological Analysis

Bacteriological characteristics were determined as described by Bezuidenhout *et al.*, (2002). The Most Probable Number multiple tube technique was used for coliform enumeration. All plates were incubated at 37°C for 24hrs. Presumptive colonies were confirmed by gram staining and biochemical reactions and each plate was given a positive or negative score. Isolates were confirmed by some conventional biochemical test SCA [7-8].

## RESULTS

The physicochemical data of the bore water samples collected in July 2013. The results of the samples vary with different collecting places because of the different nature of the soil. The physicochemical analysis (Color, odour, Turbidity, conductivity, total hardness, chlorides, fluorides, nitrites and alkalinity) is presented in Table 1. The pH of the water samples ranged from 7.7 to 7.9. while the turbidity of water samples also ranged from 1.0-2.0, total hardness of water samples ranges from 200-280 (mg/l) while With the exception of alkalinity of water samples ranges from 550-800, electrical conductivity of water samples ranges from 1400-2800 at 28°C (mmhos/cm). The chlorides in four water samples ranges from 220-820 (mg/l) fluorides present in water samples ranges from 0.6-1.0. The colour of the water samples was colorless and odour is unobjectionable. Finally nitrites present in three samples A, B, C and absence of nitrites is seen in sample D collected from bore water tiruchanoor area (Table 1).

Results of the bacteriological analysis of the water sample are presented in Table 2. The total viable counts for all water samples were quite high ranging from 6.3 x 10<sup>6</sup> cfu/ml to 2.01 x 10<sup>7</sup>cfu/ml. The most probable number (MPN) for presumptive total coliform count of the water. Samples ranged from 1,600 to >1,800 MPN per 100ml. Water samples B, and C had total Coliform count greater than 1,800 MPN per 100ml while sample D had the lowest total coliform Count of 180MPN per 100ml.

**Table 1. Physicochemical Analysis of Bore Water Samples from Water Sources present in Tirupati Rural Areas**

Parameters	Sample A Bore water (Renigunta north side)	Sample B Bore water (Renigunta south side)	Sample C Bore water (Tirupati R.T.C bus stand)	Sample D Bore water (Tiruchanoor temple)	WHO Standard	EPA Standard
PH	7.7	7.9	7.8	7.9	6.5	6.5-8.5
Conductivity (mmhos/cm)	2800	1900	1400	1800	1500	1500
Color (HU)	Colorless 5	Colorless 5	Colorless 5	Colorless 5	Colorless 5-6	Colorless 15
Odour	Un- objectionable	Un- objectionable	Un-objectionable	Un-objectionable	Un- objectionable	Un- objectionable
Turbidity (NTU)	1.0	1.0	1.0	1.0	6.0	0.5
Alkalinity (600mg/l)	800	550	650	800	600	600
Total Hardness (mg/l)	280	240	200	250	500	500
Carbonate Hardness (mg/l)	280	240	200	250	400	600
Chlorides (mg/l)	820	500	220	460	200	250
Fluorides (mg/l)	0.8	0.8	0.6	1.0	0.6	1.0-1.5
Nitrites	present	present	present	present	NS	NS

U = Unobjectionable; NS- No Standard



**Table 2. Bacteriological Analysis of Water**

Sample Code	Bore water Samples	Total Coliform count
A	Bore water 1	1,600
B	Bore water 2	>1,800
C	Bore water 3	>1,800
D	Bore water 4	180
	WHO Standard	Zero per 100ml
	EPA standard	Zero

**COLCLUSION**

In conclusion, the concentrations of the investigated major salts, carbonates and bacteria in the bore water samples from these communities in the tirupathi rural areas were found high the guidelines for drinking waters given by the World Health Organization (WHO).

Further research on other communities in this region for drinking water analyses is required as levels of contaminants may vary due to different soil types, water chemistry and different human activities. No correlations were found between metal concentrations in the drinking water samples.

**REFERENCES**

1. Dada OO, Okuofu CA and Obele E. (1990a). Fecal pollution of well water in Zaria City, Nigeria. *Savannah*, 10, 1-5
2. Dada OO, Okuofu CA and Yusuf Z. (1990b). The relationship between residual chlorine and bacteriological quality of tap water in the water distribution system of Zaria Nigeria. *Savannah*, 10 (2), 95-101.
3. Edema MO, Omemu AM and Fapetu OM. (2001). Microbiological and physicochemical analysis of different sources of drinking water. *Nigerian Journal of Microbiology*, 15, 57-61.
4. Osuinde. M.I. and Eneuzie, N.R. (1999). Bacteriological analysis of ground water. *Nigeria Journal of Microbiology*, 13, 47-54.
5. Anonymous, (1996). Guidelines for drinking water quality (2), 231,
6. World Health Organization Analysis of mine waters using X-ray fluorescence spectrometry, *Polish Journal of Environmental Studies*, 9, 429.
7. Bezuidenhout CC, Mthembu N, Puckree T and Lin J. (2002). Microbiological evaluation of the Mhlathuze River, Kwazulu-Natal (RSA). *Water SA*, 28, 281-286.
8. Standing Committee of Analysts. (2002). *The microbiology of drinking water. Part 1-Water quality and public health methods for the examination of waters and associated materials*. Environment Agency.
9. EPA, (2003). US Environmental Protection Agency Safe Drinking Water Act. EPA 816 – F – 03 –016.
10. EPA. (2002). US Environment Protection Agency, Safe Drinking Water Act Ammendment [http:// www. epa. gov/safe water /mcl. Html](http://www.epa.gov/safe-water/mcl.html)

