



# MULTIGENERATIONAL STUDY TO ACCESS THE MATERNOTOXICITY OF SLUDGE LEACHATE FROM TEXTILE INDUSTRIES AT PALI (INDIA) IN SWISS ALBINO MICE

Himani Tiwari

Water and sanitation specialist, Health of the Urban Poor Institute of Health Management Research (IIHMR), P.D.Marg, Sanganer Airport Jaipur-302011, Rajasthan, India.

Article Info	ABSTRACT
<p><i>Received 25/12/2013</i> <i>Revised 15/01/2014</i> <i>Accepted 18/01/2014</i></p> <p><b>Key words:</b> Leachate, Maternal Toxicity, Prepartum, Postpartum, Lacrimation.</p>	<p>Water and soil pollution is inherited through the uncontrolled and indiscriminate discharge of pollutants from domestic, commercial and industrial sources into water bodies. This is visible everywhere and the water pollution problem is progressively mounting in our country. An attempt has been made to compare the maternotoxic effect of sludge leachate administered during two generations. Here the leachate obtained is from CETP, Pali which receives effluent exclusively from textile and dyeing industries located at Pali. It is diluted with water according to low level exposure dose concentration of 1/1000 and then administered to Swiss albino mice along with control group receiving simple tap water for consecutive three generations in order to access the long term effect of leachate. Research finding revealed that the treated sludge leachate from CETP, Pali at the tested dose level produced maternal toxicity to certain extent at both the generations. Maternal toxicity due to sludge leachate was evident by significant reduction in body weight gain during both prepartum and postpartum period, substantial loss of body fur and restlessness in F<sub>1</sub> generation. No significant adverse effect was seen in first generation while no second generation was observed indicating the potential toxicity of leachate and ability to accumulate and persist inside the body. The toxic symptoms in mothers were muscular tremors, ataxia, convulsions, hyper salivation, lacrimation and restlessness.</p>

## INTRODUCTION

Industrialization is believed to cause inevitable problems, such as pollution of air, water and soil. Water pollution due to industrial processes has attained serious dimensions in India [1]. Both, the quality and quantity of ground water is severely threatened by industrial sewage. Among the industries, textile industry plays a major role in modern civilization. It is an important industry in Rajasthan, accounting for nearly 20 percent of the investment made in the state, contributing over 7.5 percent of India's production of cotton and blended yarn (235,000 tonnes in 2002-03) and over 5 percent of fabrics (60

millionsq meters). Rajasthan has a leading position in spinning of polyester viscose yarn & synthetic suiting and processing. State is also famous for printing & dyeing of low cost, low weight fabric. Jodhpur, Pali, Balotra, Jasol and Bituja are the major clusters of small scale industries engaged in printing and dyeing of low cost fabric. Around 1640 industries are presently operating in these clusters [2]. Water pollution due to textile industry is the topic of major concern as they discharge large quantity of effluent into nearby water bodies. [3]. Central Pollution Control Board has listed the dye industry as one of the heavily polluted industries [4]. Textile mill operations consist of weaving, dyeing, printing and finishing [5]. Many processes involve several steps, each contributing a particular type of waste, which may invite many diseases: both occupational and general [6] and consequently escalating the economic cost. Textile industry has long been known to pose health risks causing respiratory

Corresponding Author

**Dr. Himani Tiwari**

Email:- dr.himanitiwari@gmail.com



diseases like byssinosis and asthma from cotton dust exposure [7, 8] and noise induced hearing loss [9]. The above situation can be well depicted in Western Rajasthan, in India, on both the sides of river Bandi (located in Pali) that is considered as the lifeline of people living there. It houses a number of industries (textile and dyeing) that have seen a phenomenal growth during the last two decades. Studies conducted by Mohnot and Dugar [10] and Mohnot and Durve [11] have reported that various industrial units located in the three towns of Jodhpur, Pali and Balotra use about 77000 – 80000 tonnes of chemicals annually. However, potentially hazardous agents and situations are encountered in this industry, some of which might even influence the reproductive health [12]. These include solvents, dyes, noise, heat, vibrations, and prolonged standing, heavy metals etc. Humans may also be affected from exposure to agents that interfere with ovulation or spermatogenesis [13]. Exposure to such chemical compounds can produce a spectrum of adverse reproductive effects including chromosomal changes, mutations, sperm abnormalities, early or late foetal loss, still births, decreased birth weights, altered sex ratio, birth defects and childhood malignancies [14]. In the present study, an attempt has been made to study the undesired reproductive effects of chronic, low-level exposure to sludge obtained from CETP which receives effluent exclusively from textile and dyeing industries located at Pali, on reproductive parameters in Swiss Albino mice as some compounds may persistently accumulate in the body for longer period. The major part of the study involves finding out its long term teratogenic effect for continuous two generations.

## MATERIALS AND METHODS

**Study area:** The industrial effluent affected area is located along the river Bandi. In Pali town there are about 1640 industries comprising of dyeing and printing units. Approximately 36 Million Litre Per Day (mlpd) industrial effluents containing high pH, Chemical Oxygen Demand (COD), Biological Oxygen Demand (BOD), Total Dissolved Solid (TDS), Total Suspended Solid (TSS), sulphates and sodium are generated and discharged every day in the Bandi river [3,5,6,12,14,15].

**Test substance:** Test substance used is sludge, collected from drying beds of Combined Effluent Treatment Plant (CETP), Pali. The biologically and chemically treated sludge was collected, dried in oven, powdered and mixed in the ratio of 1:1. It was diluted 10 times with water and then homogenized and filtered. This filtrate served as 100% leachate, which was then diluted with water according to low low-level exposure to sludge i.e. dose concentration of 1/1000.

**Test animal:** Five to six weeks old Swiss albino mice, weighing about 20 g, were paired in the ratio of 3 females: 1 male. The females were checked for the presence of vaginal plug every morning to determine the pregnancy in

the females. The day a vaginal plug was seen was taken as day 0 of gestation and the female was presumed to be pregnant. Such females were caged singly and were given low-level exposure to sludge daily throughout in drinking water for three generations. The control animals received tap water *ad libitum* for the same period.

## Experimental protocol

Weaning animals were randomly assigned to control or test groups. The males and females in each group ( $F_0$  generation) were separately caged. The control animals received tap water and the test animals were given dose concentration of leachate 1/1000 in drinking water for 60 days. The animals in respective groups were then kept together for mating to obtain the  $F_1$  pups. The  $F_1$  pups underwent examination for any external malformations and their survival and weights were recorded till weaning. When the  $F_1$  pups attained sexual maturity, they were caged to obtain the  $F_2$  generation. The  $F_2$  pups also underwent through similar observations as the  $F_1$  pups. The data was statistically evaluated using Student's t-test [16] and one tailed Mann - Whitney U - test [17]. Females showing the presence of the vaginal plug were separated and weighed on alternate days throughout the experimental period. Maternal weight gain during the gestation period i.e. prepartum weight of the pregnant females was noted along with the post partum weight till the pups weaned. Weights of neonatal male and female pups, their survival and sex ratios were recorded at birth, and then on days 4, 7, 14, & 21 after birth [3]. The food and water consumption by the test animals was also recorded on weekly basis.

## RESULTS AND DISCUSSION

It was observed that for the first few weeks the amount of water consumed by the animals was more than their control counterparts during  $F_0$  generation but after that it reduced significantly during  $F_1$  generation. (Table 1). The experimental females of  $F_0$  parental generations did not show any changes in maternal weight gain (Table 2). The average prepartum weight gained by these dams during all the phases of gestation was less ( $12.05 \pm 2.32g$ ) as compared to the weight gained by the control females ( $15.28 \pm 2.60$ ) but no maternal mortality occurred. The postpartum weight gained by lactating females also seemed to be quite less ( $1.67 \pm 1.03$ ) than that of the control females ( $3.43 \pm 1.34$ ) but the difference was not found to be statistically significant (Table 3), when student t-test was applied to the available data. Similar relationship between maternal and fetal weight loss was observed by Hovland [18] after treating mice with Cadmium. The probable cause for such an association, studied by Himani [19] may be that as a result of the administration of leachate containing various heavy metals, organochlorines, dyes, inorganic compounds, sulphates, acids and alkalies, the maternal body are put under stress and this in turn might affect the growing fetus leading to its growth retardation and hence reduced fetal weight. The neonates of  $F_0$  generation formed the  $F_1$  generation. The litter size also didn't alter much




( $3.66 \pm 1.73$ ) from the control ( $3.77 \pm 1.63$ ) (Table 4). The surviving ability of the  $F_1$  pups was assessed from their viability index and weaning index. Weight of  $F_1$  pups was recorded on postnatal days 1, 4, 7, 14 and 21 to study their growth pattern. Neonates of dosed group revealed low viability index on postnatal days 4, 7, 14 and by the day 21 which was 79.62% as compared to 85.17% of the controls (Table 5). The observation of Growth of pups was not affected much although reduced to  $8.41 \pm 1.96$  as compared to  $10.93 \pm 2.05$  of controls. (Table 6). The  $F_1$  pups were observed for gross external malformations and their weights were taken. No stunted, immature or dead fetuses were born to any of the female of this group.

When the  $F_1$  pups attained sexual maturity, they were housed in breeding cages in the ratio of three females to one male, in order to obtain the  $F_2$  generation. Further when the  $F_1$  pups reached sexual maturity after weaning, the treated animals showed significant increase in mortality rate. The control group produced  $F_2$  generation successfully while no treated  $F_1$  animals survived to

produce  $F_2$  generation. This could be due to the toxic effect of the leachate administered, on the reproductive performance of the mice. The observations of neonates revealed low viability index, reduced growth and weaning index. It may be attributed to the reason that harmful compounds present in the dose leachate were excreted in the milk without being detoxified and thus exerted their influence and proved lethal to the pups. Such toxic effect on suckling infants feeding only on mother's milk was studied by Kavlock et al, [20]. Large number of neonatal deaths may also be attributed to diminished secretory function of mammary gland [21].

Bagnell and Ellenberger [22] reported that tetrachloroethylene, a solvent present in dry-cleaning fluid, has been reported to cause cholestatic jaundice in a breast fed infant. The mother was exposed to vapours chronically, which accumulated in her breast milk. This chlorinated hydrocarbon is therefore fetotoxic. An extensive list of solvents shown to be fetotoxic in various animal systems was tabulated by Wilson [23].

### **$F_0$ PARENTAL ANIMALS**



**Table 1. Data on water consumption from F<sub>0</sub> to F<sub>2</sub> generations of mice**

Average water consumption for two months prior to mating F <sub>0</sub> parental generation			Average water consumption by F <sub>1</sub> generation			Average water consumption by F <sub>2</sub> generation		
No. of weeks	Control	Experimental	No. of weeks	Control	Experimental	No. of Weeks	Control	Experimental
1	4.03±0.27	10.34±0.26**	1	5.27±1.17	4.70±0.31	1	5.10±0.20	6.23±0.64
2	6.41±0.41	9.49±0.78*	2	9.25±3.15	9.10±1.90	2	6.40±0.51	6.93±0.12
3	5.20±0.93	8.45±0.90	3	7.87±0.32	8.83±0.17	3	6.05±0.15	5.85±0.05
4	6.54±0.09	8.45±0.91	4	7.80±0.21	8.70±0.49	4	6.37±0.52	4.90±0.90*
5	6.33±0.26	7.79±0.50	5	7.75±0.15	9.40±0.40	5	4.53±0.17	2.60±0.30**
6	6.25±0.44	6.98±0.24	6	8.43±0.554	7.80±0.82	6	7.75±0.35	0
7	5.97±0.19	6.07±0.23	7	8.60±0.21	7.93±0.20	7	6.90±0.20	0
8	5.43±0.13	7.33±0.22**	8	8.50±0.25	7.50±0.27*	8	6.93±0.13	0
			9	7.67±0.81	7.23±0.83	9	4.53±0.37	0
			10	7.57±0.18	6.65±0.35	10	5.59±0.43	0
			11	8.33±0.22	7.10±0.10	11	6.78±0.21	0
			12	8.00±0.20	7.58±0.24	12	7.86±0.14	0
			13	7.57±0.18	6.65±0.35			
			14	7.87±0.67	6.60±0.15*			
			15	8.00±0.10	7.00±0.27			
			16	7.67±0.37	6.76±0.56			

1-8 Weeks Fo Parental gen. Given 2 months, prior mating treatment of 1/1000 leachate 8-24 Weeks Fo Parental gen. Kept together to produce F1 pups (Treatment continued) 24-35 Weeks F1 Parental gen. kept together to produce F2 pups (treatment continued) Student's t test: \*Significant difference (p<0.05) \*\* highly significant difference (p<0.01).

**Table 2. Pre Partum weight of females**

Weight Gained in (g) by Control Females During						Weight Gained in (g) By Experimental females during				
	Pre-implantation period Day 1-6	Organogenetic period Day 6-14	Growth period Day 14-22	Complete gestation period	Maternal mortality	Pre Implantation Period Day 1-6	Organogenetic Period Day 6-14	Growth Period Day 14-22	Complete Gestation Period	Maternal Mortality
F0	2.72±1.08	6.40±1.60	6.16±1.98	15.28±2.60	0	1.8±1.20	5.40±1.12	4.84±1.88	12.05±2.32	0
F1	1.88±0.61	2.26±0.61	2.1±0.66	7.32±1.32	0	0.26±0.64	0.12±0.36	0.14±0.40	0.52±0.72	0

Student's t test: \*Significant difference (p<0.05) \*\* highly significant difference (p<0.01)

**Table 3. Post partum weight of females**

Weight Gained in (g) by Control Females During					Weight Gained in (g) By Experimental females during			
	1 <sup>st</sup> post-Partum Week	2 <sup>nd</sup> post-Partum Week	3 <sup>rd</sup> post-Partum Week	3 post-Partum Weeks	1 <sup>st</sup> post-Partum Week	2 <sup>nd</sup> post-Partum Week	3 <sup>rd</sup> post-Partum Week	3 post-Partum Weeks
F0	1.05±0.99	1.73±1.03	0.85±0.86	3.43±1.34	0.72±0.67	0.73±0.68	0.6±0.73	1.67±1.03
F1	1.05±0.99	1.74±1.03	0.89±0.86	3.68±2.88	No female survived in F1 generation			

Student's t test: \*Significant difference (p<0.05) \*\* highly significant difference (p<0.01)

**Table 4. Effect of leachate on neonates of F<sub>0</sub> and F<sub>1</sub> generation**

Control group					Experimental group			
Neonates of gen.	litter size	sex ratio M:F	average pup weight at birth (gm)	immature or stunted	litter size	sex ratio m: f	Average pup weight at birth (gm)	Immature or stunted
F <sub>0</sub>	3.77±1.63	2:0	1.73±0.44	0	3.66±1.73	1:1	1.48±0.58	0
F <sub>1</sub>	3.8 ±0.33	2:1	1.49±0.60	0	1 immature, 2 resorbed. All females died during gestation period			



**Table 5. Effect of leachate on viability and weaning index of neonates**







Viability Index (%) of control pups on postnatal days						Viability Index (%) of Experimental pups on postnatal days				
Gen.	4th	7th	14th	21st	Weaning Index (%)	4th	7th	14th	21st	Weaning Index (%)
F <sub>1</sub>	96.9	95.04	85.17	85.17	85.17	93.33	72.95	72.95	72.95	79.62
F <sub>2</sub>	94.42	91.08	73.32	73.32	73.32	No pups obtained in this gen.				

Student's t test: \*Significant difference (p<0.05) \*\* highly significant difference (p<0.01)


**Table 6. Effect of leachate on growth index of pups**

Effect of leachate on growth index of pups										
Growth Index (g) of Control pups On following postnatal days						Growth Index (g) of Experimental pups On following postnatal days				
Gen.	1st	4th	7th	14th	21st	1st	4th	7th	14th	21 <sup>st</sup>
F <sub>1</sub>	1.71±0.46	2.43±0.2	3.80±0.56	6.13±1.3	10.93±2.05	1.60±0.49	2.12±0.70	3.16±1.00	4.72±1.65	8.41±1.96
F <sub>2</sub>	1.66±0.10	2.43±0.9	3.82±0.13	5.48±1.42	9.21±2.35	No pups obtained in this gen.				

Student's t test: \*Significant difference (p<0.05) \*\* highly significant difference (p<0.01)







## SUMMARY AND CONCLUSION

It can be concluded that no significant adverse effect was seen in first generation while no second generation was observed indicating the potential toxicity of leachate and ability to accumulate and persist inside the body. The present study has clearly brought out that the sludge from textile industry, which is dumped in open landfills, and with the potential of mixing with ground and surface water, may pose serious threat to human babies and fetuses and other mammalian fauna of the area. Thus our

investigation is a step towards evaluating such adverse effects on the reproductive teratological and embryological aspects.

## ACKNOWLEDGEMENTS

The author is very grateful to Rajasthan University for providing laboratory facilities for carrying out this work. The author is also grateful to Indian Institute of Health Management Research, IIHMR, and Jaipur for providing support and motivation.

## REFERENCES

- Bhosale LJ. (1985). Effect of water pollution on plants. In, Current pollution researches in India (Eds. R. K. Trivedy and P. K. Goel), Environmental. Publication, Karad, India.
- Rajasthan State Pollution Control Board, (2010). Guidelines for Abatement of Pollution In Textile Industry Published by Rajasthan State Pollution Control Board 4, Institutional Area, JhalanaDungri, Jaipur [http://210.212.96.131/rpcb/Guidelines/textile\\_guidelines\\_final.pdf](http://210.212.96.131/rpcb/Guidelines/textile_guidelines_final.pdf)
- Himani I Soni, Sharma S and Verma M. (2011). Effect of dye waste water untreated sludge from Sanganer on postnatal development of swiss albino mice. *Indian journal of Environmental sciences*, 15 (1), 81-84
- CPCB (Central pollution Control Board). (1990). Minimal national standards, Dye and dye intermediate industry document. Series, COINDS/34/1990.
- Himani. (2012). Assessment of Teratogenecity and Embryo Toxicity of Dye Wastewater Untreated Sludge from Sanganer on Swiss Albino Mice when Administered during Growth Period of Gestation. *J Health Med Inform*, 3, 111.
- Himani I. Soni PP Bakre and Bhatnagar P. (2008). Assessment of teratogenecity and embryotoxicity of sludge from textile industries at Pali (India) in Swiss albino mice exposed during organogenetic period. *J. Environ. Biol.*, 29(6), 965-969.
- BouhuysRevieByssinosis, (1978). Scheduled Asthama in the textile industry , Lung , 976/1978, Volume 154, Issue 1, 3-16
- Torén, K. (2010). Wood and Textile Industries, in Occupational and Environmental Lung Diseases (eds S. M. Tarlo, P. Cullinan and B. Nemery), John Wiley & Sons, Ltd, Chichester, UK. doi, 10.1002/9780470710425.ch17
- Md. Yusuf Haider, Ahmmad Taous, Manjur Rahim, AHM Zahurul Huq, M Abdullah. (2008). Noise induced hearing loss among the textile industry workers. *Bangladesh J of Otorhinolaryngology*, 14(2) , 39-45
- Mohnot SM and Dugar S, (1987). Textile industry and water pollution problem of western Rajasthan. In, Environmental degradation of western Rajasthan. (Eds, S.M. Mohnot and M.M. Bhandari). Jodhpur University Press. 63-71.
- Mohnot SM and Durve VS. (1989). Evaluation of the hazards of untreated and treated textile dyeing and printing waste on mammals. Technical report VIII-61. Man and Biosphere Programme. Govt. of India. New Delhi.
- Suryavathi V, Sharma S, Sharma S, Saxena P, Pandey S, Grover R, Kumar S, Sharma KP. (2005). Acute toxicity of textile dye wastewaters (untreated and treated) of Sanganer on male reproductive systems of albino rats and mice Reprod Toxicol, 19(4), 547-56.
- Olshan AF and Faustman E. (1992). Male medicated developmental toxicity. *Ann. Rev. Public. Hlth*, 14, 159- 181 (1993).
- Gupta SC. (1992). Industrialisation and chemical hazards, Swasth Hind, *Water Pollution*, 124.
- Himani Tiwari and Anil Bhargava. (2013), Maternotoxicity of Leachate from CETP Pali, When Administered to Swiss Albino Mice during Various Stages of Reproductive Cycle. *Research Journal of Chemical and Environmental Sciences*, 1(2), 42-48



16. Tiwari H. (2013). Reproductive Performance of Swiss Albino Mice Treated with Leachate from CETP Pali, Administered During Various Stages of Reproductive Cycle. *J Health Med Informat*, 4, 128.
17. Ipsen J and Feigl P. (1970). Bancroft's Introduction to Biostatistics. 2nd Edn. Harper and Row Publishers. New York, Evanston and London.
18. Sokal RR and FJ Rohlf. (1973). Introduction to Biostatistics. W.H. Freeman and Company.
19. Himani Tiwari. (2012). Assessment of Teratogenicity and Embryo Toxicity of Dye Wastewater Untreated Sludge from Sanganer on Swiss Albino Mice when Administered During Growth Period of Gestation. *Water Research & Development*, 2(1&2), 48-53.
20. Hovland DN Jr, Machado AF, Scott Jr WJ. and Collons MD. (1999). Differential sensitivity of the SWV and C57BL/6 mouse strains to the teratogenic action of single administration of cadmium given throughout the period of anterior neuropore closure. *Teratology*, 60, 13-21.
21. Kavlock RJ, Chernoff N, Hanisch RC and Gray LE. Perinatal toxicity of endrin in rodents. II. (1981). Foetotoxic effects of prenatal exposure in rats and mice. *Toxicology*, 21, 141-150.
22. Barnes JM, Denz FA. (1951). the chronic toxicity of p-nitrophenyl diethyl thiophosphate (E. 605), a long-term feeding experiment with rats. *J Hyg (Lond)* 49, 430-441.
23. Bagnell PC, Ellenberger HA. (1977). Obstructive jaundice due to a chlorinated hydrocarbon in breast milk. *CMAJ*, 117, 1047.
24. Wilson JG. (1977). Teratogenic effects of environmental chemicals. *Fed Proc*, 36, 1698-1703.

