




INTERNATIONAL JOURNAL OF NURSING EDUCATION & RESEARCH



Journal homepage: www.mcmed.us/journal/ijner

MANAGEMENT OF HIGH BLOOD CHOLESTEROL LEVELS THROUGH CUPPING THERAPY IN CLINICALLY HEALTHY YOUNG MEN

Izharul Hasan^{1*}, MdTanwir Alam², Suboohi Irshad³

¹Dept of Juris and Toxicology, A & U Tibbia College & Hospital, Karol Bagh, New Delhi, India.

²Dept of Preventive and Social Medicine, AIUMC Muzaffarnagar, UP, India.

³Dept of Gynecology and Obstetrics, A & U Tibbia College & Hospital, Karol Bagh, New Delhi, India.

Article Info

Received 25/02/2014

Revised 15/03/2014

Accepted 18/03/2014

Key words: Cupping therapy, High Blood Cholesterol.

ABSTRACT

High blood pressure is a major cause of various diseases such as stroke and high cholesterol. High cholesterol and cause a variety of deadly diseases. The aim of this study was to determine if a reduction in serum lipoproteins, especially LDL cholesterol, is a preventive approach against atherosclerosis. Phlebotomy has been a recommended method to reduce serum lipoprotein levels. As cholesterol affects the blood flow it can also impact cardiovascular fitness and sufferers may find that they get out of breath more easily or just generally feel less fit and healthy or more tired. The present study was conducted to investigate the effects of wet cupping on serum lipoprotein concentrations. In this case it is observed that cupping therapy may be an effective method of reducing LDL cholesterol in men and consequently may have a preventive effect against atherosclerosis.

INTRODUCTION

Defluoridation is removal of excess fluorides from Cholesterol is a waxy, fat-like substance that is found in all cells of the body. Hyperlipidemia refers to increased levels of lipids (fats) in the blood, including cholesterol and triglycerides. Although hyperlipidemia does not cause symptoms, it can significantly increase risk of developing cardiovascular disease, including disease of blood vessels supplying the heart (coronary artery disease), brain (cerebrovascular disease), and limbs (peripheral vascular disease). These conditions can in turn lead to chest pain, heart attacks, strokes, and other problems. Because of these risks, treatment is often recommended for people with hyperlipidemia [1, 2].

When cholesterol levels are in the normal range,

in a healthy body, the blood flows freely through the veins and arteries. When the cholesterol levels are high, it starts forming clots (plaques) in the blood vessels causing hypertension (high BP), angina (chest pain), heart attacks, strokes and peripheral vascular diseases. As cholesterol is a fat that is insoluble in blood, it needs something to carry it or transport it in our body. LDL, HDL, VLDL are small molecules called lipoproteins [3].

Blood is watery and cholesterol is fatty. Just like oil and water, the two do not mix. So, in order to travel in the bloodstream, cholesterol is carried in small packages called lipoproteins (lip-o-PRO-teens). The small packages are made of fat (lipid) on the inside and proteins on the outside. Two kinds of lipoproteins carry cholesterol throughout the body. It is important to have healthy levels of both:

LDL (Low Density Lipoprotein)

Cholesterol is sometimes called "bad" cholesterol.

Corresponding Author

IzharulHasan

Email:-drizharnium@gmail.com

Case Report



High LDL cholesterol leads to a buildup of cholesterol in arteries. The higher the LDL level in blood, the greater chance to have for getting heart disease [4].

HDL (High Density Lipoprotein)

Cholesterol is sometimes called "good" cholesterol. HDL carries cholesterol from other parts of body back to liver. The liver removes the cholesterol from body. The higher HDL cholesterol level, the lower chance of getting heart disease.

Lipid Profile

The lipid profile is a group of tests that are often ordered together to determine risk of coronary heart disease. The tests that make up a lipid profile are tests that have been shown to be good indicators of whether someone is likely to have a heart attack or stroke caused by blockage of blood vessels i.e hardening of the arteries [4].

Tests Included In a Lipid Profile

The lipid profile includes total cholesterol, HDL cholesterol (often called good cholesterol), LDL cholesterol (often called bad cholesterol), and triglycerides. The lipid profile is used to guide physicians in deciding how a person at risk should be treated. The results of the lipid profile are considered along with other known risk factors of heart disease to develop a plan of treatment and follow-up.

High Blood Cholesterol

Too much cholesterol (ko LES terol) in the blood, or high blood cholesterol, can be serious. People with high blood cholesterol have a greater chance of getting heart disease. High blood cholesterol itself does not cause symptoms; so many people are unaware that their cholesterol level is too high [4].

CAUSES OF HIGH BLOOD CHOLESTEROL [5]

Factors those can be Control

Diet

Cholesterol is found in foods that come from animal sources, such as egg yolks, meat, and cheese. Some foods have fats that raise cholesterol level. Saturated fat raises low-density lipoprotein (LDL) cholesterol level more than anything else in diet. Saturated fat is found in some meats, dairy products, chocolate, baked goods, and deep-fried and processed foods.

Trans fatty acids raise LDL cholesterol and lower high-density lipoprotein (HDL) cholesterol. Trans fats are made when hydrogen is added to vegetable oil to harden it. Trans fats are found in some fried and processed foods. Limiting foods with cholesterol, saturated fat, and Trans fats can help to control cholesterol levels.

Physical Activity and Weight

Lack of physical activity can lead to weight gain.

Being overweight tends to raise LDL level, lower HDL level, and increases total cholesterol level. (Total cholesterol is a measure of the total amount of cholesterol in the blood, including LDL and HDL.) Routine physical activity can help to lose weight and lower LDL cholesterol. Being physically active also can help to raise HDL cholesterol level.

Factors those Can't Control

Heredity

High blood cholesterol can run in families. An inherited condition called familial hypercholesterolemia external link icon causes very high LDL cholesterol. ("Inherited" means the condition is passed from parents to children through genes.) This condition begins at birth, and it may cause a heart attack at an early age.

Age and Sex

Starting at puberty, men often have lower levels of HDL cholesterol than women. As women and men age, their LDL cholesterol levels often rise. Before age 55, women usually have lower LDL cholesterol levels than men. However, after age 55, women can have higher LDL levels than men.

Signs and Symptoms of High Blood Cholesterol

High blood cholesterol usually has no signs or symptoms. Thus, many people don't know that their cholesterol levels are too high.

Diagnosis of High Blood Cholesterol [4]

High blood cholesterol is diagnosed by checking levels of cholesterol in blood. It is best to have a blood test called a lipid profile to measure cholesterol levels. Most people will need to "fast" (not eat or drink anything) for 9 to 12 hours before taking the test.

Total Cholesterol Level	Total Cholesterol Category
Less than 200 mg/dL	Desirable
200–239 mg/dL	Borderline high
240 mg/dL and higher	High

LDL Cholesterol Level	LDL Cholesterol Category
Less than 100 mg/dL	Optimal
100–129 mg/dL	Near optimal/above optimal
130–159 mg/dL	Borderline high
160–189 mg/dL	High
190 mg/dL and higher	Very high



HDL Cholesterol Level	HDL Cholesterol Category
Less than 40 mg/dL	A major risk factor for heart disease
40–59 mg/dL	The higher, the better
60 mg/dL and higher	Considered protective against heart disease

Dangers of High Cholesterol Levels

High cholesterol levels can cause:

Atherosclerosis: narrowing of the arteries [6].

Higher coronary heart disease risk: an abnormality of the arteries that supply blood and oxygen to the heart.

- Overweight and obesity
- Lack of physical activity
- Cigarette smoking
- Excessive alcohol use
- A very high carbohydrate diet
- Certain diseases and medicines
- Some genetic disorders

MODE OF TREATMENT

Cupping therapy is a very effective but under-utilized healing technique in Unani system of medicine. There are various different techniques of cupping, involving different levels of suction, heat, and duration of therapy. Stationary cups are typically placed on the patient in an area of local stagnation with a medium level of suction for 10-15 minutes. Timing and level of suction is to be adjusted depending on the type of tissue or structure which is being treated. Cupping therapy stimulates the nerves, muscles, blood vessels and subcutaneous glands and causes a series of nerve and endocrine system responses so as to improve systemic blood circulation and can eliminate blood stasis and bad cholesterol, increase permeability of blood vessels, promote blood circulations so as to improve blood supply to the kidneys and help much alleviate renal ischemia and hypoxia states.

During wet cupping, a mild suction is created using a cup that is left in place for about three to five minutes. Then removes the cup and uses a small scalpel to make superficial skin incisions. After the procedure, the site may be covered with an antibiotic ointment/betadine lotion bandage to prevent infection. The skin's appearance generally returns to normal within 6-7 days.

CASE PROFILE

MrRameez age 25 yrs, MESCO LAB Sr No. 12, lipid profile report on dated 16/11/2013. Total cholesterol 251 mg/dl, Triglycerides 540 mg/dl, VLDL 108 mg/dl, Total cholesterol /HDL ratio 5.229.

This patient was on cupping therapy only, one

Heart attack: occurs when the supply of blood and oxygen to an area of heart muscle is blocked, usually by a clot in a coronary artery. This causes your heart muscle to die [7].

Angina: chest pain or discomfort that occurs when your heart muscle does not get enough blood.

Stroke and mini stroke: occurs when a blood clot blocks an artery or vein, interrupting the flow to an area of the brain. Can also occur when a blood vessel breaks. Brain cells begin to die [8, 9].

Factors may be responsible to raise triglyceride level include:

time in 15 days, so far total 4 times therapy was done accordingly and then send for lipid profile check up in the above same LAB MESCO on dated 08/01/2014, Sr No. 20, and reports is surprising as of follows:

Informed Consent

Patients were properly informed (before treatment) about the potential for cup marks, temporary discoloration of the skin that may range from pale brown to darker red/purple depending on level of stagnation and whether the condition is of a deficient or excess, hot or cold nature. Cup marks are evidence of the removal of toxins trapped in the muscle and skin layers. Patients were also be informed that the areas of cupping should be protected from cold, wind, or water for the rest of the day after treatment.

Contraindications

Current general opinion is that cupping should not be used to treat the low back or abdomen of pregnant women, due to the possibility that through its actions on regulating the autonomic nervous system, cupping may stimulate uterine contractions, which may potentially negatively influence a high-risk pregnancy.

Preventive Measures Can Be Adopted To Reduce Cholesterol Level [10, 11]

Lifestyle

- Most people, especially those whose only risk factor has been lifestyle, can generally get their cholesterol and triglyceride levels back to normal by
- Doing plenty of exercise.
- Eating plenty of fruits, vegetables, whole grains, oats, and good quality fats.
- Avoiding foods with saturated fats.
- Getting plenty of sleep (8 hours each night).
- Bringing your bodyweight back to normal.
- Avoiding alcohol.
- Stopping smoking.



Picture 1. Patient on 1st visitPicture 2. Patient on 2nd visit

Picture 3. Patient lipid profile report before therapy

MESCO
DIAGNOSTIC CENTRE
ISO 9000 CERTIFIED DIAGNOSTIC LAB

Committed to Service

Date	16/11/2013	Srl. No.	12	Sex	M
Name	MR. RAMEEZ KHAN	Age	26 Yrs.		
Ref. By	SELF				

Test Name	Value	Unit	Normal Value
LIPID PROFILE			
TOTAL CHOLESTEROL	251.00	mg/dL	130 - 200
TRIGLYCERIDES	540.00	mg/dL	60 - 200
H D L CHOLESTEROL	48	mg/dL	30 - 70
V L D L	108	mg/dL	5.0 - 40.0
L D L CHOLESTEROL	95	mg/dL	85 - 130
TOTAL CHOLESTEROL/HDL RATIO	5.229		3.0 - 5.0
CHOLESTEROL/LDL	2.642		1.5 - 3.5
LIPID PROF INTERPRET			

**** End Of Report ****

Picture 4. patient lipid profile report after therapy

Date		Ser. No.		Sex	
08/01/2014		20		M	
Name		Age			
MR. RAMEEZ		25 Yrs.			
Ref. By					
DR. IZHAR					
Test Name	Value	Unit	Normal Value		
LIPID PROFILE					
TOTAL CHOLESTEROL	216.00	mg/dL	130 - 200		
TRIGLYCERIDES	209.00	mg/dL	60 - 200		
H D L CHOLESTEROL	52	mg/dL	30 - 70		
V L D L	41.8	mg/dL	5.0 - 40.0		
L D L CHOLESTEROL	122.2	mg/dL	85 - 130		
TOTAL CHOLESTEROL/HDL RATIO	4.154		3.0 - 5.0		
CHOLESTEROL/LDL	1.768		1.5 - 3.5		
LIPID PROF INTERPRET					
HAEMATATOLOGY					
HAEMOGLOBIN (Hb)	15.8	gm/dl	13.0 - 18.0		
**** End Of Report ****					
Checked By: 		Dr. V. JAIN MBBS MD(PATH) Consultant Pathologist			

REFERENCES

1. Third Report of the National Cholesterol Education Program (NCEP) Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) final report. *Circulation*, 2002, 106, 3143.
2. Robert S Rosenson. Patient information, High cholesterol and lipids (hyperlipidemia) Beyond the Basics <http://www.uptodate.com/contents/high-cholesterol-and-lipids-hyperlipidemia>, access on dated 07/01/2014.
3. Common queries about cholesterol, lipid profile, VLDL, HDL, triglycerides. Access online at <http://health.india.com/diseases-conditions> on dated 08/01/2014.
4. <http://www.pathologylabindia.com/patients/healthcare>, access online on dated 08/01/2014.
5. Causes High Blood Cholesterol, access online at <http://www.nhlbi.nih.gov/> on dated 08/01/2014.
6. "Hardening of the arteries" U.S. National Library of Medicine, U.S. Department of Health and Human Services National Institutes of Health. Accessed on dated 09/01/2014.
7. "Heart Attack Risk Assessment" The American Heart Association. Accessed on dated 09/01 2014.
8. "Controllable Risk Factors, Cholesterol" National Stroke Association. Accessed on dated 10/01/2014.
9. Genest J, Libby P. Lipoprotein disorders and cardiovascular disease. In, Bonow RO, Mann DL, Zipes DP, and Libby P,



- eds. Braunwald's Heart Disease, A Textbook of Cardiovascular Medicine. 9th ed. Philadelphia, Pa, *Saunders Elsevier*, 2011, chap 47.
10. Guide To Lowering Cholesterol With TLC" U.S. Department Of Health And Human Services, National Institutes of Health, National Heart, Lung, and Blood Institute. Accessed on dated 10/01/2014.
 11. <http://www.medicalnewstoday.com/> accessed on dated 10/01/2014.

